

BNL-NCS-51655 - Rev. 87
UC-34c
(Physics-Nuclear—TIC-4500)

EVALUATED NUCLEAR STRUCTURE DATA FILE

A Manual for Preparation of Data Sets

J.K. Tuli

April 1987

NATIONAL NUCLEAR DATA CENTER
BROOKHAVEN NATIONAL LABORATORY
ASSOCIATED UNIVERSITIES, INC.
UPTON, LONG ISLAND, NEW YORK 11973
UNDER CONTRACT NO. DE-AC02-76CH00016
UNITED STATES DEPARTMENT OF ENERGY

Electronic Edition by
Jay Z. James
The Isotopes Project
Lawrence Berkeley Laboratory, Berkeley, CA 94720
under USDOE Contract No. DE-AC03-76SF00098
December 1992

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency, contractor or subcontractor thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency, contractor or subcontractor thereof.

Printed in the United States of America

Available from

National Technical Information Service
U.S. Department of Commerce
5285 Port Royal Road
Springfield, VA 22161

NTIS price codes:

Printed Copy: A04; Microfiche Copy: A01

CONTENTS

CHAPTERS

I.	INTRODUCTION	1
II.	GENERAL STRUCTURE OF THE DATA FILE	
	A. General Organization	2
	B. Data Set Structure	4
	C. File Storage and Transmittal.....	5
III.	STANDARD ONE-CARD RECORD FORMATS	
	A. Introduction	6
	B. The Standard One-Card Record Formats.....	6
	C. Summary	17
IV.	RECORDS CONTAINING MORE THAN ONE CARD	
	A. Card Enumeration	19
	B. Formats for Continuation Cards.....	19
	C. Allowed Data Types on Continuation Record.....	19
V.	DETAILED FIELD DESCRIPTIONS	22
VI.	ENSDF/2 FORMAT FOR ELECTRONIC PUBLICATION	28

APPENDICES

A.	CHARACTER SET	28
B.	FORMAT FOR COMMENTS DATA SET	31
C.	SAMPLE INPUT/OUTPUT	33
D.	ENSDF TRANSLATION DICTIONARY	43
E.	DATA EVALUATION CENTERS.....	57
F.	NUCLEAR DATA SHEETS POLICIES, CONVENTIONS-----	59
	1. Language Style	60
	2. General Policies—Theory	62
	3. Bases for spin and parity assignments	64
	4. Conventions	67
	5. Abbreviations and symbols.....	68

FIGURES

1.	ENSDF Organization Chart.....	3
2.	Data Set Structure	4
3.	Format Summary	18

EVALUATED NUCLEAR STRUCTURE DATA FILE

A Manual for Preparation of Data Sets

J. K. Tuli

I. INTRODUCTION

This manual¹ describes the organization and structure of the Evaluated Nuclear Structure Data File (ENSDF). This computer-based file is maintained by the National Nuclear Data Center (NNDC) at Brookhaven National Laboratory for the international Nuclear Structure and Decay Data Network.²

For every mass number (presently, $A \leq 263$), the Evaluated Nuclear Structure Data File (ENSDF) contains evaluated structure information. For masses $A \geq 45$, this information is documented in the *Nuclear Data Sheets*; for $A < 45$, ENSDF is based on compilations published in the journal *Nuclear Physics*. The information in ENSDF is updated by mass chains with a present cycle time of approximately six years.

The author gratefully acknowledges many suggestions and comments received during the revision of this manual. Special thanks are due to M. J. Martin (ORNL), R. B. Firestone (LBL), and the following colleagues at NNDC: S. Pearlstein, M. R. Bhat, T. W. Burrows, and R. R. Kinsey. This research was supported by the Office of Basic Energy Sciences, U. S. Department of Energy.

¹The format for ENSDF was first designed by W. B. Ewbank and M. R. Schmorak at the Nuclear Data Project, Oak Ridge National Laboratory, and was described in the Rept ORNL-5054/R1 (February 1978). The present report describes the current format and supersedes both the ORNL report and the Report BNL-NCS 51655 (March 1983) by J. K. Tuli.

²Coordinated by the International Atomic Energy Agency, Vienna—see Appendix E for list of evaluation centers.

II. GENERAL ORGANIZATION AND STRUCTURE OF THE DATA FILE

A. General Organization

The Evaluated Nuclear Structure Data File (ENSDF) is made up of a collection of “data sets” which present one of the following kinds of information:

1. The evaluated results of a single experiment, e.g., a radioactive decay or a nuclear reaction.
2. The combined evaluated results of a number of experiments of the same kind, e.g., (Heavy ion, $xn\gamma$), Coulomb excitation, etc.
3. The adopted properties of the nucleus.
4. The references used in all the data sets for the given mass number. This data set is based upon reference codes (key numbers) used in various data sets for a given mass number and is added to the file by NNDC.
5. The summary information for a mass chain giving information, e.g., evaluator’s name and affiliations, cutoff date, *Nuclear Data Sheets* publication details, etc.

The data sets in ENSDF are organized by their mass number. Within a mass number the data sets are of two kinds:

- i. Data sets which contain information pertaining to the complete mass chain. These data sets contain information of the type (4) and (5) given above.
- ii. Data sets belonging to a given nucleus (Z -value)

Data sets (ii), i.e. for a given nucleus (Z -value) consist of the following:

- a. Adopted data set (only one per Z -value) giving adopted properties of the levels and radiations seen in that nucleus.
- b. Data sets giving information of the type (1) or (2) above.

If there is more than one data set of type (1) or (2) for a given nucleus, then an adopted data set is *required* for that nucleus. If there is only one data set for a given nucleus, then that set is assumed also to present the adopted properties for that nucleus.

The general organization of ENSDF is shown schematically in Figure 1.

Evaluated Nuclear Structure Data File

Organization Chart




Figure 1.

Data Set Structure




Figure 2.

B. Data Set Structure

A data set is composed of records, each record is made up of one or more 80-column card images. Data set structure is given in Figure 2 and is described below:

A data set *must* begin with an **IDENTIFICATION** record and *must* end with an **END** record (a blank card). Between these two records, there can be as many additional records as are needed to describe fully the experimental or the evaluated data.

Immediately following the **IDENTIFICATION** record is a group of records which contain information about the entire data set (#1 and #2 in Figure 2). The general **COMMENT (C)**, **NORMALIZATION (N)**, **Q-VALUE (Q)**, **PARENT (P)**, and **CROSS-REFERENCE (X)** records are of this type. Not all of these records are included in every data set. For example, **Q-VALUE (Q)** and **CROSS-REFERENCE (X)** records normally appear only in adopted data sets, while the **PARENT (P)** record is given only in radioactive decay data sets.

The body of a data set (#3 and #4 in Figure 2) is composed of numeric data records which describe the measured or deduced properties of levels, γ rays, α particles, etc. These records are associated with the level which decays (for **GAMMA** records) or the level which is populated (for **BETA**, **EC**, **ALPHA**, or **DELAYED-PARTICLE** records). Thus, each **LEVEL** record is followed by a group of records describing β , ϵ , or (delayed-) particle decay into the level and γ -ray out of the level (#4 in Figure 2). The **LEVEL** records, and the corresponding radiation records, are placed in the data set in order of increasing energy.

If a **GAMMA**, **ALPHA**, **EC**, or **BETA** record properly belongs in a data set but cannot be associated with any particular level, then the record should be placed in the data set *before any LEVEL* records (#3 in Figure 2).

The placement of **COMMENT** records is described in Chapter III.

C File Storage and Transmittal

The data sets sent to NNDC for inclusion in ENSDF can be in any order, as the file is stored in a direct access mode (by data sets) using a data base management system. Copies of the file are transmitted in the form of a sequential file on magnetic tape. The data sets in the sequential file are arranged by mass numbers in increasing numerical order. For a given mass number the data sets are organized as given in Figure 1, ordering them from left to right. Decay data sets are placed under the daughter nucleus and are ordered by A, Z and then the excitation energy of the parent nucleus. The reaction data sets are given under the residual nucleus and are ordered by the A, Z of the target nucleus followed by the A, Z of the incident particle and then by the energy of the incident particle.

III. STANDARD ONE-CARD RECORD FORMATS

A. Introduction

In most cases, all information for a record can be placed on a single 80-column card. A “standard” format has been defined for each one-card record, such that the most commonly used quantities can be placed on a single card. The standard formats are described in this section for each record. If a needed quantity is not included in the standard format or if a value will not fit within the field defined for the value by the standard format, or if a record cannot be contained on a single card, then additional cards can be prepared as described in Chapter IV (for examples, see Appendix C). Note that many of the analysis programs, at present, do not process standard fields when placed on the continuation cards.

B. The Standard One-Card Record Formats

Record formats are given below in the same order in which they would normally be encountered in a data set. Conditions under which each record may appear or be required are given in parentheses. The format descriptions give the fields (in inclusive card-column numbers), the field names (the formal “name” of the quantity that goes into the field), and a brief field description. Card columns not explicitly included in the fields are expected to be blank. A detailed description of each field can be found in the reference section noted. (Any numerical field left blank usually implies that the numerical information is lacking. Numbers will usually be assumed to be positive unless stated otherwise.) Numbers can be entered anywhere in the appropriate field (i.e., there is no need to left-adjust or right-adjust.)

1. THE IDENTIFICATION RECORD

(Required for all data sets. Must precede all other records.)

<u>Field(Col.)</u>	<u>Name</u>	<u>Description</u>	<u>Reference</u>
1-5	NUCID	Nucleus identification	V.1
6-9		Blank	
10-39	DSID	Data set identification	V.2
40-64	DSREF	References to main supporting publications and analyses	V.3
65-74	PUB	Publication information	V.4
75-80	DATE	The date (year/month/day) when the data set was placed in ENSDF (entered automatically by computer)	V.5

Note: In the rare case when the DSID field is insufficient for dataset identification, it may be continued on a second identification record with columns 1-39 defined as above, except that column 6 will contain an alphanumeric character and columns 40-80 will be blank.

2. THE Q-VALUE RECORD

(Required for adopted data sets. If there is only one data set for the nuclide, then the Q-value record should be given in that data set. Must precede L, C, B, E, A, DP records. If signs are not given, they are assumed to be positive.)

<u>Field(Col.)</u>	<u>Name</u>	<u>Description</u>	<u>Reference</u>
1-5	NUCID	Nucleus identification	V.1
6		Blank	
7		Must be blank	
8	Q	Letter ‘Q’ is required	
9		Must be blank	
10-19	Q⁻	Total energy (keV) available for β^- decay of the ground state. ($Q^- > 0$ if β^- decay is energetically possible. $Q^- < 0$ represents the Q_E energy of the Z+1 (Z = proton number) isobar.)	V.10
20-21	DQ⁻	Standard uncertainty in Q^-	V.11
22-29	SN	Neutron separation energy in keV	V.10
30-31	DSN	Standard uncertainty in SN	V.11
32-39	SP	Proton separation energy in keV	V.10
40-41	DSP	Standard uncertainty in SP	V.11
42-49	QA	Total energy (keV) available for α decay of the ground state	V.10
50-55	DQA	Standard uncertainty in QA	V.11
56-80	QREF	Reference citation(s) for the Q-values	V.3

3. THE CROSS-REFERENCE RECORD

(Given only in adopted data sets. Must precede **L, G, B, E, A, DP** records.)

<u>Field(Col.)</u>	<u>Name</u>	<u>Description</u>	<u>Reference</u>
1-5	NUCID	Nucleus identification	V.1
6		Blank	
7		Must be blank	
8	X	Letter 'X' is required.	
9	DSSYM	Any ASCII character that uniquely identifies the data set whose DSID is given in columns 10-39.	
10-39	DSID	<i>Must exactly match one of the DSIDs used.</i>	V.2
40-80		Blank	

Note: In the *Nuclear Data Sheets* the **DSID** on the first 'X' record in the data set will be identified with character 'A', and second **DSID** with 'B', and so on, irrespective of **DSSYM** on the X card. Only the first 14 **DSIDs** on 'X' records are given different symbols. All the rest are given the symbol 'O' (for others). By merely reshuffling the X records, evaluators can ascertain the **DSIDs** that will be identified individually. This has no effect on the file and affects only the published output.

4. THE COMMENT RECORD

i. General Comments

(Must precede all **L, C, B, E, A, DP** records.)

<u>Field(Col.)</u>	<u>Name</u>	<u>Description</u>	<u>Reference</u>
1-5	NUCID	Nucleus identification	V.1
6		Blank,	
7	C	Any alphanumeric character other than '1' for continuation records Letter 'C', 'D', or 'T' is required (see notes 3 – 5 below)	
8	RTYPE	Blank or type of records to which the comment pertains	V.6
9		Blank, or symbol for a particle, e.g., N, P, etc.	
10-80	CTEXT	Text of the comment [See ENSDF Translation Dictionary (Appendix D)]	V.7

NOTES:

1. The comment refers only to records of specified **RTYPE** given in that data set. The comment will normally appear only in the table for that **RTYPE** in the output. For example, if the comment is on levels ('L' in column 8) it will appear only in the level properties table.
2. If column 8 is blank, then the comment refers to the whole data set. In the case that **NUCID** contains only the mass number (columns 4-9 blank), the comment refers to the whole mass-chain A.
3. Letter 'T' in place of 'C' in column 7 of a comment record indicates to the output programs that this record should be reproduced "as is" and the blanks in the record should not be squeezed out. *See examples in Appendix C.*
4. Letter 'D' in place of 'C' in column 7 of a comment record indicates to the output programs that this is a documentation record and should be ignored. This record will also be ignored by the various analysis programs.
5. Lower case letters 'c' and 't' in column 7 of a comment record indicate to the output programs that **CTEXT** in these records should not be translated. These will appear as written in the *Nuclear Data Sheets*. In this mode one can write special characters directly, for example, "[g]" for γ , "{+238}Pu" for ^{238}Pu . See Appendix A for list of special characters.

ii. Record Comments

(Must follow the record to which the comment pertains)

<u>Field(Col.)</u>	<u>Name</u>	<u>Description</u>	<u>Reference</u>
1-5	NUCID	Nucleus identification	V.1
6		Blank,	
7	C	Any alphanumeric character other than "1" for continuation records Letter 'C' or 'D' is required. <i>See notes 4 and 5 in the section on General Comments.</i>	
8	RTYPE	Record type being commented upon	V.6
9	PSYM	Blank, or symbol for a particle, e.g., N, P, etc.	
10-80	SYM\$ or SYM,SYM,...\$	SYM = type of data being commented upon. Specified SYMs must be followed by a "\$" except as in note 1 below.	V.8
10-80	CTEXT	Text of comment follows the '\$'. On continuation comment records, CTEXT may start in col. 10, and SYM or SYMs are not repeated. [See , ENSDF Translation Dictionary. (Appendix D)]	V.7

NOTES:

1. The old format, where **SYMs** were specified in columns 10-19, will be accepted without the '\$' delimiter as long as column 19 is a blank. In this case comment text begins in column 20.
2. Record comments placed following a record of the same **RTYPE** refer only to that one record. (For example, a comment record with "CL" in columns 7-8 and "T\$" in columns 10-11 placed following the level record for the second-excited state refers only to the half-life of the second-excited state.)

iii. Footnote Comments

(Must precede **L, C, B, E, A, DP** records)

<u>Field</u>	<u>Name</u>	<u>Description</u>	<u>Reference</u>
1-9		same as in ii (record comments)	
10-80	SYM\$ or SYM,SYM,...\$ or SYM(FLAG) \$ or SYM(FLAG). SYM(FLAG),...\$	V.8 SYM = see note 1 below; FLAG = any ASCII alphanumeric character or string of alphanumeric characters <i>Field must end with a "\$"</i> <i>See note 1 on Record comments for exception</i>	
10-80	CTEXT	Text of comment follows "\$" On continuation comment records SYM or SYM(FLAG) are not repeated. [See ENSDF Translation Dictionary (Appendix D)]	V.7

NOTES:

1. **SYM** can be only one of the following:
 - The fields defined in formatted **L, C, B, E, A, DP** records.
 - TITLE**—see note 2 below. **FLAG** with this **SYM** is not allowed.
 - BAND**—this **SYM** must be accompanied by a **FLAG**. Note also that text following '\$' delimiter, or in columns 20-80 in old format, will appear as the band label in the drawing. Any other information on that band should, therefore, be given on continuation records.
 - CONF**—same as **BAND** except that **FLAG** is optional. Without the given configuration comment will apply to all levels.**FLAG**
2. Footnote without FLAG
 - This refers to all records of the specified **RTYPE** in the data set.
 - The footnote will normally appear only in the table for that **RTYPE** in the output. For example, if the footnote is on levels ('L' in column 8) it will appear only in the level properties table.
 - If the word "TITLE" was used as **SYM** the footnote will apply to the table heading.
3. Footnote with FLAG
 - Only those records are footnoted for which footnote flags are given, see note 4 below.
 - Only those data values of data types specified by **SYM** which is associated with a given **FLAG** are footnoted.

4. Footnote **FLAG** must be either a single character placed in column 77 of the formatted record or a string of characters assigned to a special data type called FLAG on the following continuation record.

Examples of flags on a continuation record:

```
152EU2 G FLAG=ABCD$
156GD2 L FLAG=KMP$
```

5. No footnotes are allowed for records of **RTYPE : N, P, or Q**.
6. To change the standard label heading of a formatted field, e.g. S to C²S for L records, **CTEXT** should have the form LABEL=name, where "name" is the new label desired. The new label should be kept as short as possible. Note that **FLAG** cannot be specified with relabeling; also any other comment on the relabelled field must appear on a different record.

Examples of field relabel:

```
156GD CL S$LABEL=C2S
156GD CL S$LABEL=DSIGMA/DOMEGA (45 DEG)
```

5. THE PARENT RECORD

(Required for all decay data sets, except *IT* and *SF* decays. Must precede *L, C, B, E, A, DP* records.)

<u>Field(Col.)</u>	<u>Name</u>	<u>Description</u>	<u>Reference</u>
1-5	NUCID	Parent nucleus identification	V.1
6		Blank	
7		Must be blank	
8	P	Letter 'P' is required	
9		Blank	
10-19	E	Energy of the decaying level in keV (0 for ground state)	V.18
20-21	DE	Standard uncertainty in E	V.11
22-39	J	Spin and parity	V.20
40-49	T	Half-life; units <i>must</i> be given	V.14
50-55	DT	Standard uncertainty in T	V.12
56-64		Must be blank	
65-74	QP	Ground-state Q-value in keV (total energy available for g.s.-g.s. transition); it will always be a positive number	V.9
75-76	DQP	Standard uncertainty in QP	V.11
77-80		Must be blank	

Note: More than one parent card is allowed in a data set. If the decay scheme is due to more than one parent level then separate **P** records should be given for each parent level.

6. THE NORMALIZATION RECORD

(Must precede **L, C, B, E, A, DP** records

. Required if an absolute normalization is possible ;used mainly with decay and (n, γ) reaction data sets.)

<u>Field(Col)</u>	<u>Name</u>	<u>Description</u>	<u>Reference</u>
1-5	NUCID	Nucleus (daughter/product) identificatio	V.1
6		Blank	
7		Must be blank	
8	N	Letter 'N' is required	
9		Blank	
10-19	NR	Multiplier for converting relative <i>photon</i> intensity RI (in the GAMMA record) to <i>photons</i> per 100 decays of the parent through the decay branch or to photons per 100 neutron captures in an (n, γ) reaction. <i>Required</i> if the absolute photon intensity can be calculated.	V.9
20-21	DNR	Standard uncertainty in NR	V.11
22-29	NT	Multiplier for converting relative transition intensity (including conversion electrons) [TI in the GAMMA record] to transitions per 100 decays of the parent through this decay branch or per 100 neutron captures in an (n, γ) reaction. <i>Required</i> if TI are given in the GAMMA record and the normalization is known.	V.9
30-31	DNT	standard uncertainty in NT	V.11
32-39	BR	Branching ratio multiplier for converting intensity per 100 decays through this decay branch to intensity per 100 decays of the parent nucleus. <i>Required if known.</i>	V.9
40-41	DBR	Standard uncertainty in BR	V.11
42-49	NB[†]	Multiplier for converting relative β^- and ϵ intensities (IB in the B- record; IB, IE, TI in the EC record) to intensities per 100 decays through this decay branch. <i>Required if known.</i>	V.9
50-55	DNB	Standard uncertainty in NB	V.11
56-62	NP	Multiplier for converting per hundred delayed-transition intensities to per hundred decays of precursor	V.9
63-64	DNP	standard uncertainty in NP	V.11
65-80		Must be blank	

[†]Note: Normally β^- and ϵ intensities are given as per 100 parent decays and therefore NB=1.0. If a value other than 1.0 is given it should be remembered that the multiplier for conversion to per 100 decays is NB \times BR. Also, the uncertainties in I(β^-) will be calculated from addition of three quantities $\Delta(I(\beta^-))$, DBR and DNB in quadrature. Unless the uncertainties are precisely known it is recommended that NB be given without uncertainty.

6A. The Production Normalization RECORD

(Must follow N record, if N record present. It should be given when G records present>)

<u>Field (Col.)</u>	<u>Name</u>	<u>Description</u>	<u>Reference</u>
1-5	NUCID	Nucleus (Daughter/Product) identification	V.1
6		Blank	
7	P	Letter "P" (for production) is required	
8	N	Letter "N" is required	
9		Must be blank	
10-19	NR×BR	Multiplier for converting relative photon intensity (RI in the GAMMA record) to <i>photons</i> per 100 decays of the parent. (Normally NR×BR). If left blank (NR-DNR) × (BR DBR) from N record will be used for normalization.	V.9
20-21	UNC[†]	Standard uncertainty in NR×BR	V.11
22-29	NT*BR	Multiplier for converting relative <i>transition</i> intensity (including [†] conversion electrons) [TI in the GAMMA record] to <i>transitions</i> per 100 decays of the parent. (Normally NT×BR) If left blank (NT DNT) × (BR DBR) from N record will be used for normalization.	V.9
30-31	UNC[†]	Standard uncertainty in NT×BR	V.11
42-49	NB × BR	Multiplier for converting relative β^- and ϵ intensities (IB in the B– record; IB, IE, TI in the EC record) to intensities per 100 decays. If left blank (NT DNT) × (BR DBR) from N record will be used for normalization.	V.9
50-55	UNC[†]	Standard uncertainty in (NB DNT) × (BR DBR)	V.11
56-62	NP	Same as in "N" record	
63-64	UNC[†]	Standard uncertainty in NP	V.11
77	COM	Blank or "C" (for comment) If blank, comment associated with the intensity option will appear in the drawing in the <i>Nuclear Data Sheets</i> . If letter "C" is given, the desired comment to appear in the drawing should be given on the continuation ("nPN") record(s), col. 10-80.	
78	OPT	Intensity Option. Option as to what intensity to display in the drawings in the <i>Nuclear Data Sheets</i> . The available options are given below (default option 3).	

<u>Option number</u>	<u>Intensity displayed</u>	<u>Comment in drawing</u>
1	TI or RI(1+ α)	Relative I(γ +ce)
2	TI × NT or RI × NR × (1+ α)	I(γ +ce) per 100 (mode) decays
3	TI × NT × BR or RI × BR × NR (1+ α)	I(γ +ce) per 100 parent decays
4	RI × NR × BR	I(γ) per 100 parent decays
5	RI	Relative (γ)
6	RI	Relative photon branching from each level
7	RI	% photon branching from each level

[†] If left blank no uncertainty will appear in the publication.

7. THE LEVEL RECORD

(Optional, although a data set usually has at least one.)

<u>Field(Col.)</u>	<u>Name</u>	<u>Description</u>	<u>Reference</u>
1-5	NUCID	Nucleus identification	V.1
6		Blank Any alphanumeric character other than '1' for continuation records	
7		Must be blank	
8	L	Letter 'L' is required	
9		Must be blank	
10-19	E	Level energy in keV — <i>Must not be blank</i>	V.18
20-21	DE	Standard uncertainty in E	V.11
22-39	J	Spin and parity	V.20
40-49	T	Half-life of the level; units <i>must</i> be given. Mean-life expressed as the width of a level, in units of energy, may also be used.	V.14
50-55	DT	Standard uncertainty in T	V.12
56-64	L	Angular momentum transfer in the reaction determining the data set. (Whether it is L_n , L_p , ΔL , etc., is determined from the DSID field of the IDENTIFICATION record.)	V.22
65-74	S	Spectroscopic strength for this level as determined from the reaction in the IDENTIFICATION record. (Spectroscopic factor for particle-exchange reactions; β for inelastic scattering.) Note: If a quantity other than spectroscopic factor is given in this field, a footnote relabelling the field is required.	V.21
75-76	DS	Standard uncertainty in S	V.11
77	C	Comment FLAG used to refer to a particular comment record	V.8
78-79	MS	Metastable state is denoted by 'M' or "M1" for the first (lowest energy) isomer; "M2", for the second isomer, etc.	V.17
80	Q	The character '?' denotes an uncertain or questionable level Letter 'S' denotes neutron or proton separation energy or a level expected but not observed.	

8. THE BETA (β^-) RECORD

(Must follow the **LEVEL** record for the level which is fed by the β^- .)

<u>Field(Col.)</u>	<u>Name</u>	<u>Description</u>	<u>Reference</u>
1-5	NUCID	Nucleus identification	V.1
6		Blank Any alphanumeric character other than '1' for continuation records	
7		Must be blank	
8	B	Letter "B" is required	
9		Must be blank	
10-19	E	Endpoint energy of the β^- in keV <i>Given only if measured</i>	V.18
20-21	DE	Standard uncertainty in E	V.11
22-29	IB	Intensity of the β^- -decay branch [†]	V.13
30-31	DIB	Standard uncertainty in IB	V.11
42-49	LOGFT	The log <i>ft</i> for the β^- transition for uniqueness given in col. 78-79	V.9
50-55	DFT	Standard uncertainty in LOGFT	V.12
56-76		Must be blank	
77	C	Comment FLAG ('C' denotes coincidence with a following radiation . A "?" denotes probable coincidence with a following radiation.)	V.8
78-79	UN	Uniqueness classification for the β^- decay, e.g., "1U", "2U" (A blank signifies an allowed or a nonunique forbidden transition)	V.16
80	Q	The character '?' denotes an uncertain or questionable β^- decay Letter "S" denotes an expected or predicted transition	

[†] The intensity units are defined by the **Normalization** record.

9. THE EC (or EC + β^+) RECORD

(Must follow the *LEVEL* record for the level being populated in the decay.)

<u>Field(Col.)</u>	<u>Name</u>	<u>Description</u>	<u>Reference</u>
1-5	NUCID	Nucleus identification	V.1
6		Blank	
		Any alphanumeric character other than '1' for continuation records	
7		Blank	
8	E	Letter 'E' is required.	
9		Must be blank	
10-19	E	Energy for <i>electron capture</i> to the level <i>Given only if measured</i>	V.18
20-21	DE	Standard uncertainty in E	V.11
22-29	IB	Intensity of β^+ -decay branch [†]	V.13
30-31	DIB	Standard uncertainty in I	V.11
32-39	IE	Intensity of electron capture branch [†]	V.13
40-41	DIE	Standard uncertainty in IE	V.11
42-49	LOGFT	The log <i>ft</i> for ($\epsilon + \beta^+$) transition for uniqueness given in columns 78-79	V.9
50-55	DFT	Standard uncertainty in LOGFT	V.12
65-74	TI	Total ($\epsilon + \beta^+$) decay intensity [†]	V.13
75-76	DTI	Standard uncertainty in TI	V.11
77	C	Comment FLAG (Letter 'C' denotes coincidence with a following radiation. A '?' denotes probable coincidence with a following radiation.)	V.8
78-79	UN	Uniqueness classification for ϵ , β^+ decay, e.g., "IU", "2U". (A blank signifies an allowed or a nonunique forbidden transition)	V.16
80	Q	The character "?" denotes an uncertain or questionable ϵ , β^+ branch. Letter "S" denotes an expected or predicted transition	

10. THE ALPHA RECORD

(Must follow the *LEVEL* record for the level being populated in the decay.)

<u>Field(Col.)</u>	<u>Name</u>	<u>Description</u>	<u>Reference</u>
1-5	NUCID	Nucleus identification	V.1
6-		Blank	
7		Must be blank	
8	A	Letter 'A' is required	
9		Must be blank	
10-19	E	Alpha energy in keV	V.18
20-21	DE	Standard uncertainty in E	V.11
22-29	IA	Intensity of α -decay branch in <i>percent</i> of the total α decay	V.13
30-31	DIA	Standard uncertainty in IA	V.11
32-39	HF	Hindrance factor for α decay	V.9
40-41	DHF	Standard uncertainty in HF	V.12
42-76		Must be blank	
77	C	Comment FLAG (Letter 'C' denotes coincidence with a following radiation; A "?" denotes probable coincidence with a following radiation.)	V.8
78-79		Must be blank	
80	Q	The character '?' denotes uncertain or questionable α branch Letter 'S' denotes an expected or predicted α branch	

11. THE (DELAYED-) PARTICLE RECORD

(Must follow the **LEVEL** record for the level which is fed by the particle.*

Records for particles which are unassigned in a level scheme should precede the first level of the data set.)

<u>Field(Col.)</u>	<u>Name</u>	<u>Description</u>	<u>Reference</u>
1-5	NUCID	Nucleus identification	V.1
6		Blank Any alphanumeric character other than '1' for continuation records	
7		Must be blank	
8	D	Blank for prompt -, Letter 'D' for delayed-particle emission	
9	Particle	The symbol for the (delayed) particle (N=neutron, P=proton, A=alpha particle) is required	
10-19	E	Energy of the particle in keV	V.18
20-21	DE	Standard uncertainty in E	V.11
22-29	IP	Intensity of (delayed) particles in <i>percent</i> of the total (delayed-) particle emissions †	V.13
30-31	DIP	Standard uncertainty in IP	V.11
32-39	EI	Energy of the level in the "intermediate" (mass=A+1 for n, p) nucleus in case of delayed particle	V.13
40-49	T	Width of the transition in keV	V.14
50-55	DT	Uncertainty in T	V.10
56-64	L	Angular-momentum transfer of the emitted particle	V.22
65-76		Blank	
77	C	Comment FLAG used to refer to a particular comment record	V.8
78	COIN	Letter "C" denotes placement confirmed by coincidence. Symbol "?" denotes probable	V.15
79		Blank	
80	Q	The character '?' denotes an uncertain placement of the transition in the level scheme Letter "S" denotes an expected, but as yet unobserved, transition.	

* The delayed-particle record will appear in a delayed-particle data set (e.g. B-N DECAY, etc.) which should be given under the A-chain for the final nucleus. For example, "95RB B-N DECAY" should be given as data set for ⁹⁴Sr.

† The intensity units are defined by the **Normalization** record.

12. THE GAMMA RECORD

(Must follow the **LEVEL** record for the level from which the γ -ray decays. Records for γ -rays which are unassigned in a level scheme should precede the first level of the data set.)

<u>Field(Col.)</u>	<u>Name</u>	<u>Description</u>	<u>Reference</u>
1-5	NUCID	Nucleus identification	V.1
6		Blank	
		Any alphanumeric character other than '1' for continuation records	
7		Must be blank	
8	G	Letter 'G' is required	
9		Blank	
10-19	E	Energy of the γ - ray in keV – <i>must not be blank</i>	V.18
20-21	DE	Standard uncertainty in E	V.11
22-29	RI	Relative <i>photon</i> intensity [†]	V.13
30-31	DRI	Standard uncertainty in RI	V.11
32-41	M	Multipolarity of transition	V.19
42-49	MR	Mixing ratio, δ (Sign must be shown explicitly if known. If no sign is given, it will be assumed to be unknown.)	V.10
50-55	DMR	Standard uncertainty in MR	V.12
56-62	CC	Total conversion coefficient	V.9
63-64	DCC	Standard uncertainty in CC	V.11
65-74	TI	Relative total transition intensity [†]	V.13
75-76	DTI	Standard uncertainty in TI	V.11
77	C	Comment FLAG used to refer to a particular comment record. The symbol “*” denotes a multiply- placed γ -ray. The symbol “&” denotes a multiply-placed transition with <u>intensity not</u> divided. The symbol “@”denotes a multiply-placed transition with intensity suitably divided.	V.8
78	COIN	Letter “C”denotes placement confirmed by coincidence. Symbol “?”denotes questionable coincidence.	V.15
79		Blank	
80	Q	The character ‘?’ denotes an uncertain placement of the transition in the level scheme Letter”S” [†] denotes an expected , but as yet unobserved, transition.	

[†] The intensity units are defined by the **NORMALIZATION** record.

13. THE REFERENCE RECORD

(Record can occur only in Reference data set. N N DC provides the Reference data.set)

<u>Field(Col.)</u>	<u>Name</u>	<u>Description</u>	<u>Reference</u>
1-3	MASS	Mass number	
4-7		Must be blank	
8	R	Letter "R" is required	
9		Must be blank	
10-15	KEYNUM	Reference key number	V.3
16-80	REFERENCE	Abbreviated reference (from NSR file)	

14. THE END RECORD

(Required for all data sets. Must be the last record in a data set.)

<u>Field(Col.)</u>	<u>Description</u>
1-80	All columns are blank

C. Summary

The following figure (Figure 3) summarizes the standard one-card formats for all allowed record types.

IV. RECORDS CONTAINING MORE THAN ONE CARD

A. Card Enumeration

If all the information for a given record type cannot be contained on a single card, it is possible to use additional cards to describe the record fully. The first card of a record will have a blank in column 6. Subsequent cards will have characters different from blank or 1 (usually running numbers: 2 to 9 or letters A to Z).

B. Format for Continuation Cards

THE CONTINUATION RECORD

(Must follow the record of the same *RTYPE*.)

<u>Field(Col.)</u>	<u>Name</u>	<u>Description</u>
1-5	NUCID	Nucleus identification
6		Any alphanumeric character other than 1. Note: "S" is reserved for computer-produced records which will usually be suppressed in the <i>Nuclear Data Sheets</i> .
7		Must be blank
8	RTYPE	Letter corresponding to the record type L , B , E , or G
9		Must be blank
10-80	Data	<quant><op><value>[<op><value>][<ref>]\$...

In the description of **Data** above the following abbreviations have been used:

- <quant>: Standard symbol for a quantity as defined in **IV.3** below.
Notes: 1. Ratios of more than two quantities should be indicated by colons not slashes (e.g., K:L1:L2:L3, not K/L1/L2/L3).
2. See **V.23** for description of <value> when <quant>=XREF
- <op>: =, <, >, EQ, AP, LT, LE, GT, GE
Note: The last 6 operators require blanks before and after them are required.
- <value>: Numeric value with units as needed and optional uncertainty. Uncertainty is as defined in Sections **V.11** and **V.12**.
Note: For ranges, uncertainties should not be included.
To specify a bounded range of values a second operator (note that =, EQ, AP are not valid) and value are required.
See examples below.
- []: Optional.
- <ref>: 6 character key numbers, **KEYNUM** (see **V.3**), separated by commas and enclosed within parentheses, e.g., (76TU01,81BO01).
- \$: Delimiter (end of record is also a delimiter; thus '\$' should not be the last character in a record)

Examples:

```
126TE 2 G BE2W=25.3 7(70LAZM)
126I 2 L %EC+%B+=56.3 20 (77JA04)$%B- EQ 43.7 20 (77JA04)
126SN 2 B EAV=2030 60
126TE 2 L G LE 0.19 GT 0.1 (81SH15)$MOME2 AP -0.20$BE2=0.478 12
```

C. Allowed Data Types on Continuation Records

Each record type is permitted to contain only a limited (but extendable) set of data types. For example, a **GAMMA** record is not allowed to contain information of data type **DTYPE = J** (nuclear spin). Neither may a **LEVEL** record contain **LOGFT** information.

No continuations are allowed for **Q**, **N**, and **P** records. For **A** and **DP** records only data type **FLAG** is allowed on the continuation records. For the special format for the continuation of **IDENTIFICATION** records see III.B-1. The allowed record types for **LEVEL**, **GAMMA**, **B-**, and **EC** records are described below.

1. The LEVEL Record

Allowed data types **E, DE, J, T, DT, L, S, DS, C, MS, and Q** are described with the standard formats in Section III.B.6. Additional allowed data types are:

<u>TYPE</u>	<u>Description</u>
%EC, %B+, %EC+%B+, %B-, %IT, %SF, %A, %P, %N %B-N; %B-xN; ...	Percent decay of the level by ϵ , β^+ , $\epsilon+\beta^+$, β^- isomeric transition, spontaneous fission, α , proton, or neutron decay Percent delayed decay through n,xn emission. Similarly for other particle emissions, e.g., p, xp, α , x α , etc., following β^- , β^+ , or ϵ decays. <i>Note: Decay modes must be given on "2 L" card in adopted set.</i>
G	g-factor of the level
MOME1, MOME2, ...	Electric moments: dipole, quadrupole, ...
MOMM1, MOMM2, ...	Magnetic moments: dipole, quadrupole, ...
CONF	Nuclear configuration of the level
BE1, BE2, ...	Reduced electric transition probability (<i>upward</i>), given in units $e^2 \times (\text{barns})^L$, where $L = 1, 2, \dots$ for the transition from the ground state to this level
B2, B3,	2^L -pole ($L=2,3,\dots$) nuclear deformation parameter
ISPIN	Isobaric spin
ISPINZ	Z-component of isobaric spin
WIDTH,WIDTHG,WIDTHG0, WIDTHN,WIDTHP,WIDTHA	Level width, Γ , Partial- γ , $-\gamma_0$, $-n$, $-p$, $-\alpha$ widths, $\Gamma(\gamma)$, $\Gamma(\gamma_0)$, $\Gamma(n)$, $\Gamma(p)$, $\Gamma(\alpha)$, respectively
XREF	Cross-reference to other data sets for that nuclide
FLAG	Additional footnote symbols

2. The GAMMA Record

Allowed data types, **E, DE, RI, DRI, M, MR, DMR, CC, DCC, TI, DTI, C, COIN, Q** are described with the standard formats in Section III.B.7.

Additional allowed data types are:

<u>DTYPE</u>	<u>Description</u>
BE1, BE2, ...	Reduced electric transition probability (<i>downward</i>) given in units of $e^2 \times (\text{barns})^L$, where $L = 1, 2, \dots$
BE1W, BE2W, ...	Reduced electric transition probability (<i>downward</i>) given in single-particle (Weisskopf) units
BM1, BM2, ...	Reduced magnetic transition probability (<i>downward</i>) given in units of $\mu^2 \text{ N} \times (\text{barns})^{L-1}$, where $L = 1, 2, \dots$
BM1W, BM2W, ...	Reduced magnetic transition probability (<i>downward</i>) given in single-particle (Weisskopf) units
CEK, CEL, CEL1, ...	Conversion-electron (ce) intensity for K, L, L_1, \dots conversion
ECC	Measured total conversion coefficient
KC, LC, LIC,	Theoretical K-, L-, L_1 -, ... conversion coefficient
EKC, ELC, EL1C, ...	Measured K-, L-, L_1 -, ... conversion coefficient
K/L, M/L, L1/L2, ...	Conversion-electron intensity ratios
K/T, L/T, ...	Ratio of K, L, ... ce- intensity to total ($\gamma + \text{ce}$) intensity
CE	Total conversion electron intensity
FL	Final level energy
FLAG	Additional footnote symbols

3. The BETA (β^-) Record

Allowed data types **E, DE, IB, DIB, LOGFT, DFT, C, UN, Q** are described with the standard formats in Section III.B.9 Additional allowed data types are:

<u>DTYPE</u>	<u>Description</u>
EAV	Average energy of the β^- -spectrum
FLAG	Additional footnote symbols (Note: 'C' and '?' may not be used – see III.B.8 for their special meaning)

4. The EC Record

Allowed data types, **E, DE, IB, DIB, IE, DIE, LOGFT, DFT, TI, DTI, C, UN, Q** are described with the standard formats in Section III.B.9. Additional allowed data types are:

<u>DTYPE</u>	<u>Description</u>
EAV	Average energy of the β^+ spectrum
CK, CL, CM, ..., CL+	Calculated fraction of decay by electron capture from the K, L, M, ..., L+M+... shells (K, ...L, M, ..., L+ are also acceptable as DTYPE and have the same meaning.)
ECK, ECL, ECM, ..., ECL+	Measured fraction of decay by electron capture from the K, L, M, ..., L+M+... shells
CK/T, CL/T, ...	Ratio of K, L, ...ε- intensity to total ϵ intensity
FLAG	Additional footnote symbols (Note: "C" and "?" may not be used -- see III.B.9 for their special meaning)

V. DETAILED FIELD DESCRIPTIONS

1. NUCID

The standard nucleus identification consists of two parts—mass number in columns (1-3), right justified, and element name (or Z-100 for $Z > 103$) in columns 4-5, left justified. The nucleus identification must be contained *within* the field defined for it (columns 1-5). The nucleus identification *must* be included on every **IDENTIFICATION** record. It must also be included on every card of a data set except the **END** record. Comments and reference data sets pertaining to the whole A-(mass) chain evaluation contain only the A-value in the **NUCID** field.

2. DSID

The Data Set **ID** for an ENSDF data set must serve as a unique, computer-recognizable identification for the data set. For that purpose, the following rules should be strictly observed for ENSDF entries. Single blanks have meaning and should be used according to the formats below. A colon may be used to define a sub-topic. All characters must be confined to the 30 spaces allowed. *Optional fields are given in italics*. General categories are given in upper and lower cases and further defined.

GENERAL ID'S

REFERENCES
COMMENTS (see Appendix B for format for this data set)
ADOPTED LEVELS
ADOPTED LEVELS, GAMMAS

DECAY DATA SET ID'S

Parent Mode Decay (*Half - life*)

Parent should be the parent isotope symbol (e.g.) 52CR.

Mode may be one of **B+**, **B-**, **EC**, **IT**, **A**, **P**, **B-N**, **ECP**, or **SF**.

Half-life can be of the form **T** defined in **V.14.1**.

MUONIC ATOM

REACTION DATA SET ID'S

Target(Reaction), (*Reaction*), *Target(Reaction) E=Energy Qualifier*
COULOMB EXCITATION (*Reaction*)
INELASTIC SCATTERING
(HI,XNG)

Target should be the target (isotope or element) symbol.

Reaction should be a reaction symbol (e.g.) **N,P**.

Energy may be one of the following:

NUM, NUM Units (for definition of NUM see **V.9**)
NUM-NUM Units
THERMAL or TH
RESONANCE or RES

Qualifier may be one of the following RES, IAR or IAS.

EXAMPLES:

187RE B- DECAY	187OS IT DECAY (231 US)
187AU P DECAY:?	190PT A DECAY (6E11 Y)
95RB B-N DECAY	186OS(N,G) E=THERMAL
186W(N,G) E=TH: SECONDARY G'S	RE(N,N'):TOF
186W(N,G) E=RES: AVG	189OS(P,T) E=19 MEV
187OS(D,D') E=12, 17 MEV	185RE(A,2NG) E=23-42.8 MEV
187RE(D,2NG), 187RE(P,NG)	44CA(P,G) E=856, 906 KEV IAR

3. DSREF, KEYNUM, QREF

The **DSREF** and **QREF** fields may include up to three key numbers (**KEYNUM**), each of which refers to a particular publication. Additional key numbers may be placed in **COMMENT** records. *Key numbers must be left-justified and separated by commas with no blanks between the comma and the reference.* A reference key number must be of the form **YYAABB** where **YY** is a two-digit integer, **AA** are two alphabetic characters and **BB** is either a two-digit integer or consists of two alphabetic characters. Examples: **81TU01**, **81TUXY**, etc.

4. PUB

Publication information consists of the year of the A-chain publication in *Nuclear Data Sheets* denoted by a two-digit year indicator followed by the three-character code **NDS**. This may optionally be followed by a comma and other updating information, e.g., the initials of the person modifying the data set after its publication. Example: **78NDS.TWB** or **81NDS**.

5. DATE

This field is of the form **YYMMDD** where **YY**, **MM** and **DD** are two digit integers within the following ranges: **00** ≤ **YY** ≤ **99**, **01** ≤ **MM** ≤ **12**, and **01** ≤ **DD** ≤ **31**.

6. RTYPE

RTYPE is a one-letter code in column 8 that gives a name to the **RECORD** type.

<u>RTYPE</u>	<u>Description</u>
blank	May be IDENTIFICATION , general COMMENT , or END record
N	NORMALIZATION record
P	PARENT record
Q	Q-VALUE record
L	LEVEL record
G	GAMMA record
B	BETA (B⁻) record
E	EC (or EC + B⁺) record
A	ALPHA record
R	REFERENCE record
X	CROSS-REFERENCE record
DP	DELAYED PARTICLE record, or PARTICLE (column 8=blank) record. Particle symbol (e.g., "P" for proton) is given in column 9.

7. CTEXT

This field consists of free text. The various expressions used in **CTEXT** can be translated via dictionary lookup. The translation dictionary is given in Appendix D. The unit expression used in translation is the string of characters between adjacent "delimiters". The characters presently used as "delimiters" are:

b(blank) ,(comma) .b(period followed by a blank); : () - = + < > / and \$

In some cases the dictionary lookup programs look beyond the next delimiter for proper translation.

8. SYM(FLAG)

The **SYM(FLAG)** field (with **FLAG** given) is valid only for records with **RTYPE L, G, B, E, A, and DP**. However, **SYM** (without **FLAG**) may additionally be used for record types **N, P, and Q**.

FLAG can be a string of characters *optionally* separated by commas. Any character other than a comma and parenthesis can be used as a **FLAG** symbol. For **B** and **E** records, "C" cannot be used for a **FLAG**, as "C" in column 77 of **B, E, and A** records denotes coincidence. Similarly "*" and "&" for **G** records are reserved to denote multiple placement in the level/decay scheme. See notes on **SYM** and **FLAG** on page 9. Note also that **FLAG** can be used only with **BAND** and **CONF** in addition to those **SYMs** which are valid data types on a formatted card. *In fact, for BAND FLAG must be given.*

Allowed symbols to be used as **SYM** for various **RTYPE** are given below:

RTYPE	Allowed SYM
L	E, DE, J, T, DT, L, S, DS, BAND, CONF, BE1, BE2, ..., B2, B3, ..., G, ISPIN, ISPINZ, MOME1, MOME2, ..., MOMM1, MOMM2, ..., %EC+%B+, %X (X=B-, B+, EC, IT, SF, A, P, or N), %XY (X=B-, B+, or EC; Y=N, P, A, ..., or KN, KP, KA, ... where K=2,3, ..., or 9), WIDTH, WIDTHG, WIDTHG0, WIDTHN, TITLE
G	E, DE, RI, DRI, M, MR, DMR, CC, DCC, TI, DTI, BE1, BE2, BE1W, BE2W, ..., BM1, BM2, ..., BM1W, BM2W, ..., CC, CEK, CEL, CEL1, ..., KC, LC, L1C, ..., CE, EKC, ELC, EL1C, ..., K/L, M/L, L1/L2, ..., K/T, L/T, ..., TITLE
B	E, DE, IB, DIB, LOGFT, DFT, EAV, TITLE
E	E, DE, IB, DIB, IE, DIE, LOGFT, DFT, TI, DTI, EAV, CK, CL, CM, CN+, ECK, ECL, ECM, ECN+, CK/T, CL/T, TITLE
A	E, DE, IA, DIA, HF, DHF, TITLE
N	NR, DNR, NT, DNT, BR, DBR, NB, DNB
P	E, DE, J, T, DT, QP, DQP
Q	Q-, DQ- SN, DSN, SP, DSP, QA, DQA
DP	E, DE, IP, DIP, EL, T, DT, L

9. CC, NR, NT, BR, NB, QP, LOGFT, HF

These fields consist of either a blank or a single unsigned number (NUM) in one of the following forms:

1. An integer (e.g., 345)
2. A real number (e.g., 345.23)
3. An integer followed by an integer exponent (e.g., 345E- 4, 4E+5)
4. A real number followed by an integer exponent (e.g., 345.E- 4)

Note: It is desirable to write a number as '0.345' rather than '.345'. However, even if the leading '0' were omitted, presumably to save space, it will appear with a leading '0' in the *Nuclear Data Sheets*.

10. Q-, SN, SP, QA, MR

These fields have the same form as the quantities in V.9. above with the difference that they are allowed to have signature (positive or negative), and R can be within parentheses.

11. DNR, DNT, DBR, DNB, DQP, DQ-, DSN, DSP, DQA, DE, DRI, DTI, DIB, DIE, DCC, DIA, DS, DHF

These one- or two-character fields represent uncertainty in the "standard" form in the given quantity. The "standard" numeric uncertainty denotes an uncertainty in the last significant figure(s), for example, NR=0.873, DNR=11 represent a normalization factor of 0.873 ± 0.011 . Similarly QP=2.3E6, DQP=10 stand for a Q-value of $(2.3 \pm 1.0) \times 10^6$ (see also Appendix F-1). The non-numeric uncertainty, e.g., <, >, or ≥, etc. is denoted by expressions LT, GT, and GE, etc.

The allowed forms for these fields are summarized below:

1. Blank
2. An integer ≤ 99 , preferably ≤ 25 , (left or right justified)
3. One of the following expressions:

LT, GT, LE, GE, AP, CA, SY

for less than, greater than, less than or equal to, greater than or equal to, approximately equal to, calculated, and from systematics, respectively.

12. DFT, DMR, DT

These fields allow for the specification of “standard” asymmetric uncertainty. For example, $T=4.2 \text{ S DT}=+8-10$, represent a half-life $=4.2^{+0.8}_{-1.0}$ s; similarly, $MR= -3 \text{ DMR}= +1-4$ represent mixing ratio $= -3^{+1}_{-4}$ meaning a range from -7 to -2 . (Note: asymmetric uncertainties add algebraically.) When the $+/-$ construction is missing from this field, the digits or the expressions given in this field represent either the numeric “standard” symmetric or the non-numeric uncertainty as described in **V.11.** above.

To summarize this field, there are two cases:

1. Symmetric uncertainty—the field consists of an integer number or an expression of the type described in **K.** above.
2. Asymmetric uncertainty—the field is of the form $+x-y$, where x and y are integers.

13. RI, TI, IB, IE, IA

The following numbers/expressions are valid for these fields:

1. NUM (number as defined in **V.9** above)
2. (NUM)

Note: Parentheses denote that the number given has been deduced (not directly measured) or taken from other experiment(s).

14. T

The field for half-life **T** must have one of the following forms:

1. NUM–Blank--Units (i.e. number as defined in **V.9** above followed by a blank and its units). Valid symbols for units are: Y, D, H, M, S, MS, US, NS, PS, FS, AS, EV, KEV, and MEV for year, day, hour, minute, second,(s), 10^{-3} s, 10^{-6} s, 10^{-9} s, 10^{-12} s, 10^{-15} s, 10^{-18} s, eV, 10^3 eV, and 10^6 eV, respectively.
2. Word “STABLE”

Note: A question mark following half-life denotes that the assignment to that level is not certain. A comment should be given to explain the exact meaning intended.

15. COIN

This one character field can either be blank or have character “C” or ‘?’ . The character “C” denotes coincidence, while ‘?’ denotes questionable coincidence.

16. UN

This two-character field can either be blank or have an integer between 1 and 9 followed by the character “U”.

17. MS

This two-character field can either be blank or have an “M” followed by a blank or a digit between 1 and 9.

18. E

An energy field **E** can have only one of the following forms:

- 1 NUM (as defined in **V.9.** above)
- 2 NUM+A or A+NUM, where A=X, Y, Z, U, V, or W used in this order; i.e., for the first occurrence an 'X' is used, for its second occurrence a 'Y' is used, and so on.
- 3 SN+NUM, SP+NUM, SN+A, or SP+A (where A is as defined in 2. above)
- 4 A (as defined in 2. above)

Note: Parentheses are allowed for this field. They denote that the number given has been deduced (not directly measured) or taken from other experiment(s). Explanation as to what is intended should be given.

19. M

The multipolarity field can be one of the following:

1. Mult
2. Mult+Mult
3. Mult, Mult
4. NOT Mult
5. IF Mult or [MULT]

where Mult = E_L or M_L , (where L, L' are single digits – $L \geq 0, L' \geq 1$) or
 $M_{L'+E_L}$ or
 $E_{L+M_{L'}}$ or
D or Q

Note: Parentheses in the multipolarity field denote that the assignment is probable and not definite. Square brackets indicate assumed or derived assignment.

20. J

The spin-parity field can have only one of the following forms:

1. JPI
2. JPI OR JPI (' , ' (comma) can be used in place of 'OR')
3. JPI AND JPI (' & ' (ampersand) can be used in place of 'AND')
4. OP JPI (where OP is AP, LE, or GE
Note: This will be interpreted as $\pi=PI$ and J is OP J
Example: LE 5+ means $\pi=+$ and $J \leq$
5. NOT JPI
6. JPI TO JPI (' : ' (colon) can be used in place of 'TO')

Note: If parity is given in the range it will be interpreted as follows:

- a. J to J'PI means $J \leq J' \leq J'$ and $\pi=PI$
- b. JPI to J'PI' means $JPI, J=J+1 PI=\pm, \dots, J=J'-1PI=\pm, J'PI'$
- c. JPI to J' means $JPI, J=J+1 PI=\pm, \dots, J=J'-1PI=\pm, J'PI=\pm$

Examples:

- a. 3 to 6- means $J\pi=3-, 4-, 5-, 6-$,
- b. 3+ to 6- means $J\pi=3+, 4\pm, 5\pm, 6-$
- c. 3+ to 6 means $J\pi=3+, 4\pm, 5\pm, 6\pm$

7. NATURAL/UNNATURAL

In the above, $J = N$ or $N/2$ (N is a positive integer or zero)
 $PI = +$ or $-$
 $JPI = J$ or PI or J followed by PI .

Notes:

1. Parentheses in the J^π field indicate that the parenthesized value(s) is (are) based upon weak arguments. See "Bases for Spin and Parity Assignments" — Appendix F-4. Note that $JPI = (3,4)-$ is interpreted as $J=(3)$ or (4) and $\pi=-$.
2. As far as possible, do not give more than three JPI values.
3. The ranges such as 3- to 5+ are better written as 3-,4,5+.
4. Square brackets around J^π value indicate an assumed value.

21. S

This field may contain no more than three S-values, in the form of NUM defined in **V.9**, separated by a '+' or a comma, for corresponding L-values given in the L-field (columns 65-74). Parentheses are allowed and will be interpreted to mean probable values.

22. L

This field may contain no more than three integer numbers optionally preceded by LE or GE and separated by a '+' or a comma. Parentheses are allowed and will be interpreted to mean probable values. Square brackets indicate assumed or derived values.

For certain reactions the excitation L value may be accompanied by its electric or magnetic character in the form similar to multipolarity (V.19).

23. Cross Reference

The cross referencing of a record (currently allowed only for the "L" record in an ADOPTED data set) is done through specification on the continuation record and it takes the following forms:

1. NUCID 2 L XREF=ABC\$
Above record indicates that the adopted level (specified by the preceding 'L' record) has been seen in data sets "A", "B", and "C", and that the corresponding levels are unambiguous.
2. NUCID 2 L XREF=A(E1)B(E2)C(E3)\$
This record indicates that the adopted level is the same as the E1 level in data set "A" the E2 level in data set "B" etc.
3. NUCID 2 L XREF=A(E1,E2)B(E3)\$
This record indicates that the adopted level is either the E1 or the E2 level in data set "A", the E3 level in data set "B."
4. NUCID 2 L XREF=A(*E1)B(E2)\$
This record indicates that a level with energy E1 in data set "A" is associated with two adopted levels. An "*" must appear on all occurrences of a multiply assigned level. Alternatively, the notation A(*) may be used if the energy is apparent.
5. NUCID 2 L XREF=+\$
This record indicates that the adopted level has been seen in all data sets.
6. NUCID 2 L XREF=- (AB)\$
This record indicates that the adopted level has been seen in all data sets except the data sets "A" and "B".

Note: The symbols A, B, C relating to specific data sets must be defined through Cross-Reference records (see **III.B.3**).

VI. ENSDF/2 Format For Electronic Publication

Richard B. Firestone
Isotopes Project, Lawrence Berkeley National Laboratory, Berkeley CA 94720

The ENSDF/2 format for the Evaluated Nuclear Structure Data File (ENSDF) is being developed at the Lawrence Berkeley Laboratory to support the development of electronic publication of nuclear data on desktop computers. This effort has been divided into two phases. In the first phase, the ENSDF file will be enhanced to include statistical information on each dataset, explicit level, transition, and cross-reference indexing, updating of mass related data, and standardization of comment formats. This phase is designed to be an ENSDF upgrade and will not change the content of the pre-existing ENSDF file. Phase I will be completed at LBL without the requirement of evaluator assistance. The second phase of development will be the incorporation of ENSDF/2 into a data base management system which will be provided with the ENSDF desktop publication system. An evaluator shell will be provided to support interactive entry of nuclear data on the desktop. Phase II will contain internal format checking procedures and standard ENSDF calculation utilities, obviating the need for pre-existing software. The Phase I format will become the standard transmission format for ENSDF/2. It is anticipated that conventional ENSDF format data will continue to be prepared by some evaluators, although this format will not be recommended. We will provide software to convert between ENSDF and ENSDF/2 (Phases I and II).

Phase I: Software for ENSDF/2 development has been derived, in part, from codes developed to produce the 8th edition of the *Table of Isotopes*. A preliminary version of ENSDF/2 is used by the beta version of the VuENSDF program. Figure 4 summarizes the formats adopted for ENSDF/2.

Statistical Information: Interactive generation of tables and figures is facilitated by the inclusion of "I" records, following the DSID record, that indicate the quantities of level and transitions in each dataset, which fields are utilized, and other information that can improve the generation of output.

Indexing: In order to accurately support the genetic information implied by a level scheme it is necessary that the initial and final levels associated with all transitions are explicitly included in the file. We have developed a two-character index which is used to uniquely name each level in a dataset. Redundant fields 4-5 on each level record have been utilized to store this index. For γ -ray records the initial level index is inferred from the previous level record, but the final level must be calculated from available data. We do this by comparing possible final levels on the basis of energy difference, multipolarity versus ΔJ^π , +X information, and band assignment. These criteria are weighted and the final level is selected by software. Possible problems with final level selection are documented for off-line checking. The final level index is stored in fields 4-5 of the γ -ray record.

Rotational band information is also indexed. Each band comment record is assigned a single-character index which is stored in column 3 of the comment record and column 3 of the associated level records. On γ -ray records the band assignment of the initial level is stored in column 3 and that of the final level in column 2. For all records uncertainty information is moved from column 80 to column 1, and for levels isomeric information is moved from column 78 to column 2. This accommodates a significant decrease in the size of the ENSDF/2 file.

Beta and alpha-decay records have well defined final levels and their initial levels are referred to the parent record. This is not completely satisfactory because parent records are not always consistent with the associated adopted levels. This problem is solved by matching the parent record to the associated adopted level (see cross-referencing below) and storing the associated level index in columns 78-79 of the parent record and columns 4-5 of the beta or alpha record. For delayed particle data the situation is more complex because parent, level, and final level indices are all required. For delayed-particle records the final level index will be stored in columns 2-3. For beta, alpha, and delayed-particle

ENSDF/2

RECORD TYPE

SUMMARY OF ENSDF/2 FORMATS

	1	2-3	4-5	6	7	8	9	10-19	20-21	22-29	30-31	32-39	40-41	42-49	50-55	56-62	63-64	65-74	75-76	77	78	79	80
IDENT	<NUCID->							*	-DSID-				><	-REF-			><	-PUB><		-DATE-			
STATISTICS	AAA		ZZZ			I		NL	NG		NA		NB		NPAR	AZP1	AZP2		NBD				
BAND COM.		F						BAND(F)\$<				-COMMENT-											
BAND COM.		G						BAND(G)\$<				-COMMENT-											
GEN. COM.	IDC			#	C	L	A	CTEXT-															
FLAG COM.	IDC	&		#	C	L	A	CTEXT-															
XREF							X	A DSID-															
NORMALIZ.							N	A	<-NR->	<DNR>	<-NT->	<DNT>	<-BR->	<DBR>	<-NB->	<DNB>	NR*BR	DNB	NT*BR	DNB			
PARENT	<AZPAR->						P	A	<-E->	<DE->	<-JPI->	<-T->	<DT->	<-QP93	DQP>	<-QP->	<DQP>					IDP	
LEVEL	Q	M	F	IDL			L	A	<-E->	<DE->	<-JPI->	<-T->	<DT->	<-L->	<-S->	<-DS->						C	IDA
GAMMA	Q	F	G	IDL2			G	A	<-E->	<DE->	<-RI->	<DRI>	<-M->	<-MR->	<-DMR->	<CC->	<DCC>	<-TI->	<DTI>				C
BETA	Q	UNQ	IDP				B	A	<-E->	<DE->	<-IB->	<DIB>		<LOGFT>	<DFT->	FT93	DFT						
EC	Q	UNQ	IDP				E	A	<-E->	<DE->	<-IB->	<DIB>	<-IE->	<DIE>	<LOGFT>	<DFT->	FT93	DFT					
ALPHA	Q	L	IDP				A	A	<-E->	<DE->	<-IA->	<DIA>	<-HF->	<DHF>		HF93	DHF						
PARTICLE	Q	IDD	IDP				D	+	<-E->	<DE->	<-IP->	<DIP>	<-ED->	<WIDTH>	<DW->	<-L->							

comment continuation number & Flag symbol + particle symbol

Record Format Changes:

- General: Most NUCID deleted, column 9 used to designate dataset where blank=adopted, Q moved to column 1.
- Identification: * in column 10 indicates ENSDF/2 format
- Statistics: Integer a,z; number of levels, gammas, alphas, betas, parents, AZ of parents, number of bands.
- Band
 - comments: Band flag in column 3. FLAG= record used to replace comment field.
- General, Flag
 - comments: standard comment number column 1-3, flag in column 4, no more than one flag per comment.
- Normalization: Unused columns for normalization products.
- Parent: Most recent Q-value and adopted level ID added
- Level: Isomer fields moved; band assignment field, level ID added, and Adopted level ID added if not same.
- Gamma: Initial/final band flag fields and final level ID added
- Beta, EC,Alpha: UN moved, parent ID added, logft(HF) from new Q-value added.
- Particle: Parent and daughter ID's added.

records information in column 80 will be moved to column 1.

Cross-references: Cross-indexing of levels in datasets to Adopted levels is indicated, in ENSDF, by XREF records specifying the dataset, but not the specific level within that dataset, which is matched. In order to make the cross-referencing explicit, the matching adopted level ID from columns 4-5 is stored in columns 78-79 of the corresponding dataset records. This matching is performed on the basis of the existing XREF record information, and a weighted comparison of energies, J^π values, half-lives, and γ -ray deexcitation patterns. When XREF records have not been entered into the file, they can be created automatically.

Mass Data: Masses in ENSDF generally correspond to values from the Wapstra and Audi mass tables at the time of the evaluation. These values are supplemented in ENSDF/2 with the newest available values (currently Nuclear Physics **A564**, 1 (1993)). The masses are included on a "2 Q" record following the original Q record in the adopted levels dataset. In addition, the previously unused parent record fields 56-64 are used to store the newest Q-values. Recalculated $\log ft$ values or hindrance factors are stored in the unused fields 56-64 on A, E, or B records.

Comment Records: Comments in ENSDF are generally interpreted from a context sensitive dictionary. An alternative ASCII character set is also defined for ENSDF which allows a more exact description of the output text. For ENSDF/2 comments will be converted to the ASCII character set. This will change the content of character records, but not their presentation at output. An exact translation of ENSDF will require some proofreading because of the imprecise nature of the ENSDF dictionary.

ENSDF Cleanup: In ENSDF/2 the records will be reorganized in a more regular and predictable manner. This will include grouping records of a common type so that primary records are followed by secondary records and then by comments. Level records, and gamma records following each level, will be organized by energy in ascending order. Flagged comments will be given unique flag characters and included only on "FLAG=" records. Column 77 flags will also be moved to "FLAG=" records. Comment and reference information included on secondary data records will be removed to comment records. DSID information on X records will be exactly matched to actual dataset names.

Phase II: The second phase of ENSDF/2 development will divide the Phase I file into separate databases of levels, transitions, and comments. These databases will be interconnected by common indices and it will be possible to reconstruct the Phase I format database, or even original ENSDF, from these databases. Development of Phase II will be coupled to the selection of database management software to support data retrieval and display. This will allow us to access and evaluate nuclear data by property as well as by isotopic dataset. Several modifications of ENSDF/2 are anticipated in Phase II that will make the file more versatile.

Document Comments: Comments will be treated as editable documents which can be prepared in a conventional word-processor environment. Standard comments will be available for use by the evaluator, however it will be possible to create more complex comments with mathematical expressions, imbedded drawings, and a full range of available fonts and characters. This will allow the display of spectra and figures directly taken from published papers.

Reference Links: Reference keynumbers in ENSDF will be hypertext linked to the Nuclear Structure Reference file (NSR). This will require distribution of the NSR file with ENSDF/2 for these links to become effective. These hypertext links provide direct indexing of the references from the comments so that they can be retrieved and displayed with the comment information.

Numerical Data: At this time ENSDF is essentially a text file. However, it is often desirable to search the file quantitatively. We will add numerical fields to ENSDF to provide searchable values of energies

and intensities. In addition, the spins and multipolarities will be numerically searchable.

Data Entry: Phase II ENSDF/2 will have database modification capabilities which allow the user or evaluator to interactively enter or change data. These will be driven primarily by full screen table editors that will allow users to edit standard level and transition tables. Input format checking will assure the integrity of the file and help the user correctly perform level-transition indexing and cross-referencing. Application software for calculating conversion coefficients, $\log ft$ values, alpha hindrance factors, least-squares level energy calculations, decay scheme balancing and normalization, and many other utilities will be provided.

Standardization: The ENSDF/2 file will adopt standard data presentation formats which will be selected in concert with the editors and evaluators of the file. These formats will be enforced by data checking algorithms. The purpose of this standardization will be to provide users with a more standard presentation of information than was possible in the past.

APPENDIX A

CHARACTER SET

The character set available includes all characters currently used at the NNDC and provides for future expansion. The base character set is the standard seven-bit ASCII character set up to octal 173. Characters with octal values of 173 and greater are used as control characters. An alternate character set is defined which consists primarily of the Greek alphabet and special symbols. The backslash character (octal 134) is interpreted as a backspace command. An alternate character in the input file consists of two characters, a control character and the standard character equivalent of the alternate character. All available alternate characters and their standard equivalents are given in Table IV.

There are four control characters: “|” (octal 174), “~” (octal 170), “{“ (octal 173) and “}” (octal 175). The vertical bar and the tilde are used to shift the next character into the first and second alternate character sets respectively. Entire strings of characters may also be modified from their standard form. In this case the string to be modified is enclosed by the open and close brace control characters. The character immediately following the open brace is interpreted as a control character. The available control character values and their meanings are given in Table V. Modified character strings may be nested, but one may not terminate a string without terminating all strings nested within that string. The control character may be either upper or lower case .

EXAMPLES

g	will be displayed	γ
{B{+238}Pu}	will be displayed	238Pu

Special compound characters can be generated as follows:

v̄	from	n"backslash character" "
ñ	from	h"backslash character" `
λ	from	l"backslash character" `

TABLE A-1

STRING CONTROL CHARACTERS

1 OR	–	first alternate character set
2 OR ~	–	second alternate character set
+	–	superscript
–	–	subscript
I	–	italic
S	–	script
B	–	boldface
U	–	underline
O	–	overscore
E	–	elevate characters
R	–	raise base line

Note: + and – are mutually exclusive, as are I and S.

APPENDIX A (cont.)

TABLE A-2
AVAILABLE CHARACTERS

<u>OCTAL</u>	<u>STANDARD</u>	<u>ALTERNATE</u>	<u>OCTAL</u>	<u>STANDARD</u>	<u>ALTERNATE</u>
40	(blank)	(blank)*	116	N	N
41	!	©	117	O	O
42	"	—	120	P	Π
43	#	§	121	Q	Θ
44	\$	e	122	R	P
45	%	√	123	S	Σ
46	&	≡	124	T	T
47	'	°	125	U	Υ
50	(←	126	V	V
51)	→	127	W	Ω
52	*	×	130	X	Ξ
53	+	±	131	Y	Ψ
54	,	1/2	132	Z	Z
55	-	+	133	[{
56	.	∞	134		
57	/	÷	135]	}
60	0	(136	^	↑
61	1)	137	˘	↓
62	2	[140	˘	˘
63	3]	141	a	α
64	4	<	142	b	β
65	5	>	143	c	γ
66	6	√	144	d	δ
67	7	∫	145	e	ε
70	8	∏	146	f	φ
71	9	∑	147	g	γ
72	:	†	150	h	χ
73	;	‡	151	i	ι
74	<	≤	152	j	ε
75	=	≠	153	k	κ
76	>	≥	154	l	λ
77	?	≈	155	m	μ
100	@	∞ •	156	n	ν
101	A	A	157	o	ο
102	B	B	160	p	π
103	C	H	161	q	θ
104	D	Δ	162	r	ρ
105	E	E	163	s	σ
106	F	Φ	164	t	τ
107	G	Γ	165	u	υ
110	H	X	166	v	ϑ
111	I	I	167	w	ω
112	J	~	170	x	ξ
113	K	K	171	y	ψ
114	L	Λ	172	z	ζ
115	M	M			

* The first ALTERNATE blank is an invariant character, one half the width of the STANDARD blank; ; and the second ALTERNATE blank is one quarter the width of a STANDARD blank.

APPENDIX B
FORMAT FOR COMMENTS DATA SET

This data set consists only of general comment records (defined in III.B(4)). The format of the comment records is similar to general comments in other data sets except that the NUCID field will contain only the mass number, AAA, and that a SYM field is required as in a flagged comment. As in the flagged comments, the SYM field will either occupy columns 10 to 19 with column 19 being blank or the SYM will be followed by a "\$". Continuation records for a given comment are allowed with the additional feature that a new line will be started if the continuation character in column 6 is a "#" and that a new paragraph will be started if the character is a "@". This feature is intended to facilitate the entry of information into the COMM comments.

<u>SYM</u>	<u>Meaning</u>
TITL	Title of evaluation . Required if the evaluation spans several masses.
AUTH	Authors, a list of authors from the institution given in the following INST comment. A letter or number in parentheses following an author's last name will signal a permanent address which is different from that of the institution. (See PERM.)
INST	Institution, name and address of the authors' institution. The INST comment must follow the appropriate AUTH comment. The # continuation character is used so the address does not run together into one line. More than one set of AUTH and INST comments can be given if more than one institution is involved.
ABST	Abstract, should be terse and to the point. Additional details should be given under COMM comments.
CUT	Cutoff date and associated comments.
COMM	General comments on techniques used in the evaluation or on other information common to many of the isotopes.
ACKN	Acknowledgments.
PERM(a)	Permanent address of an author—the letter or number 'a' within the parentheses corresponds to the letter or number within the parentheses which follows the author's last name in the AUTH comment.
FUND	Funding—an acknowledgment of funding which will result in a footnote being added to the title.
CIT	Citation. To be added by the NDS production staff so that the publication can be correctly cited by persons using a retrieval of the A chain. The authors may leave it out.

APPENDIX C

EXAMPLES OF INPUT/OUTPUT IN ENSDF/Nuclear Data Sheets

Appendix C-1 EXAMPLE – COMMENTS DATA SET

Appendix C-2 EXAMPLE – ADOPTED LEVELS

Appendix C-3 EXAMPLE – DECAY DATA SET

APPENDIX C-- 1

Example of a Comments Data Set

```

156      COMMENTS
156  C  TITL      NUCLEAR DATA SHEETS FOR A=156
156  C  AUTH      R. G. HELMER
156  C  INST$IDAHO NATIONAL ENGINEERING LABORATORY
156 #C  EG&G IDAHO, INC.
156 #C  IDAHO FALLS, IDAHO 83415  USA
156  C  ABST      THE EXPERIMENTAL RESULTS FROM THE VARIOUS REACTION AND
156 2C  DECAY STUDIES LEADING TO NUCLIDES IN THE A=156 MASS CHAIN, AND
156 3C  ALPHA DECAYS FROM IT, HAVE BEEN REVIEWED.  THESE DATA ARE
156 4C  SUMMARIZED AND PRESENTED, TOGETHER WITH ADOPTED LEVEL SCHEMES
156 5C  AND PROPERTIES.
156  C  CUT      DATA AVAILABLE PRIOR TO JUNE 1985 HAVE BEEN EVALUATED.
156  C  ACKN      THE EVALUATOR WISHES TO THANK M. A. LEE AND C. W.
156 2C  REICH FOR MANY HELPFUL DISCUSSIONS AND A. L. FREEMAN FOR THE DATA
156 3C  ENTRY AND PROCESSING.
156  C  FUND      RESEARCH SPONSORED BY THE U. S. DEPARTMENT OF ENERGY.
156  c  COMM$In this evaluation, the following expression was used to define
156 2c  the rotational-band parameters A and B:
156 #c
156 #c      E(J) = E{-0} + AJ(J+1) + B[J(J+1)]{+2}.
156 #c
156 #c  For the levels in {+156}Gd the parameters A{-2} and A{-4}  representing
156 7c  the shift between the odd- and even-spin levels have also been
156 8c  calculated for some bands from the expressions
156 #c
156 #c      E(J) = E{-0} + AJ(J+1) - (-1){+J}A{-2}J(J+1)    for K|p= 1+
156 #c  and
156 #c      E(J) = E{-0} + A[J(J+1)-K{+2}] + B[J(J+1)-K{+2}]{+2} +
156 Dc  (-1){+J}A{-4}(J-1)J(J+1)(J+2)  for K|p= 2+.
156 #c
156 #c  In the determination of the values of these parameters, the energy
156 Gc  spacings of only the lowest levels, and minimum number of levels, were
156 Hc  used.
156 @c  The term "half-life" is used to refer to a ground state or an isomer
156 2c  for which there is a separate data set. The term "lifetime" is used to
156 3c  refer to the same quantity for any other level.
156 @c  In each group of data sets for a given element, the following customs
156 Bc  are usually maintained. All of the information concerning the
156 Cc  gamma-ray multipolarities are considered and summarized in the Adopted
156 Dc  data set, and these adopted values are also quoted in all the other
156 Ec  data sets. For level lifetimes, the data in the individual data sets
156 Fc  included only values from that type of experiment. The values which
156 Gc  summarize all of the measurements are only given in the Adopted data
156 Hc  set. The J|P values for the levels are also treated in the latter
156 Ic  way, except in so far as the authors of the experimental papers have
156 Jc  already considered the results of other experiments. Since a B(E2)
156 Kc  value to a 2+ excited state from 0+ g.s. is equivalent to the
156 Lc  lifetime of E2 G in the opposite direction, a T{-1/2} value can be
156 Mc  computed from B(E2) value. This has not been done. All T{-1/2} values
156 Nc  come only from experiments measuring this quantity.
156 @c  The ENSDF file (the computer data base from which these Data Sheets
156 9c  are produced), contains some information that is not printed in these
156 9c  Data Sheets. This includes the theoretical internal-conversion
156 9c  coefficients for each shell, where the values are significant, for
156 9c  each |g for which a multipolarity is given in the Data Sheets. Also, a
156 9c  short comment is made about the experimental methods for each
156 9c  reference. This information would be available if a copy of the ENSDF
156 9c  file were obtained.
156  C  CIT      NDS 49,383 1986

```

APPENDIX C-1 (cont'd)

Output example for Comments Data Set

Nuclear Data Sheets for A = 156*

R. G. HELMER

Idaho National Engineering Laboratory

EG&G Idaho, Inc.

Idaho Falls, Idaho 83415 USA

(Received October 10, 1985; Revised March 27, 1986)

Abstract: The experimental results from the various reaction and decay studies leading to nuclides in the A=156 mass chain and α decays from it have been reviewed. These data are summarized and presented together with adopted level schemes and properties.

Cutoff Date: Data available prior to June 1985 have been evaluated.

Organization of Material: See page v.

Acknowledgements: The evaluator wishes to thank M. A. Lee and C. W. Reich for many helpful discussions and A. L. Freeman for the data entry and processing.

General Comments: In this evaluation, the following expression was used to define the rotational-band parameters A and B:

$$E(J) = E_0 + AJ(J+1) + BJ(J+).$$

For the levels in ^{156}Gd the parameters A_2 and A_4 representing the shift between the odd- and even-spin levels have also been calculated for some bands from the expressions

$$E(J) = E_0 + AJ(J+1) - (-1)^J A_2 J(J+1) \text{ for } K\pi = 1^+$$

and

$$E(J) = E_0 + A[J(J+1) - K^2] + B[J(J+1) - K^2]^2 + (-1)^J A_4 (J-1)J(J+1)(J+2) \text{ for } K\pi = 2+.$$

In the determination of the values of these parameters the energy spacing of only the lowest levels and minimum number of levels were used.

The term half-life is used to refer to a ground state or an isomer for which there is a separate data set. The term lifetime is used to refer to the same quantity for any other level.

In each group of data sets for a given element the following customs are usually maintained. All of the information concerning the gamma-ray multiplicities are considered and summarized in the Adopted data set, and these adopted values are also quoted in all the other data sets. For level lifetimes the data in the individual data sets included only values from that type of experiment. The values which summarize all of the measurements are only given in the Adopted data set. The $J\pi$ values for the levels are also treated in the latter way except in so far as the authors of the experimental papers have already considered the results of other experiments. Since a B(E2) value to a 2+ excited state from 0+ g.s. is equivalent to the lifetime of E2 G in the opposite direction, a $T_{1/2}$ value can be computed from B(E2) value. This has not been done. All $T_{1/2}$ values come only from experiments measuring this quantity.

The ENSDF file (the computer data base from which these Data Sheets are produced) contains some information that is not printed in these Data Sheets. This includes the theoretical internal-conversion coefficients for each shell where the values are significant for each γ for which a multipolarity is given in the Data Sheets. Also a short comment is made about the experimental methods for each reference. This information would be available if a copy of the ENSDF file were obtained.

* Research sponsored by the U. S. Department of Energy.

APPENDIX C-2

Example of an Adopted Levels Data Set

```

154SM ADOPTED LEVELS 870129
154SM Q -717.0 21 7967.9 10 9086 16 -1197.2 25 85WA02
154SM D Other reactions: 154SM(P,P'G), 154SM(D,D'), and 152SM(t,p).
154SM T Summary of Adopted T Values
154SM T
154SM T E(level) Adopted Comments
154SM T 0.0 Stable
154SM T 81.9 3.02 NS 4 Weighted average of 3.03 NS 5 (67W006) and
154SM2T 3.00 NS 6 (68RI09). Other: 2.74 NS 24 (59BI10).
154SM T 266.7 172 PS 4 Weighted average of 173 PS 5 (72DI06) and
154SM2T 169 PS 10 (80J008).
154SM T 543.9 22.7 PS 6 Weighted average of 23.3 PS 7 (72DI06) and
154SM2T 22.7 PS 6 (average of two values in 80J008).
154SM T 902.8 5.93 PS 25 Weighted average of 6.2 PS 6(72DI06), 6.0 PS 4
154SM2T (77KE06), and 5.6 PS 4 (80J008).
154SM T 921.5 26 FS 4 Calculated from level width of 0.0176 eV 24
154SM T
154SM T Summary of JPI Assignments
154SM T
154SM t E(level) Adopted b- decay (n,n'g) Coulomb excit
154SM T -----
154SM T 0.0 0+ 0+ 0+ 0+
154SM T 81.9 2+ 2+ 2+ 2+
154SM T 266.7 4+ 4+ 4+ 4+
154SM T 543.9 6+ 6+ 6+ 6+
154SM T
154SM C The data on the level energies and JPI values are
154SM2C primarily from the 154PM B- DECAY.
154SM CL E From least-squares fit to G energies.
154SM CL BAND(A) KPI=0+ ground-state band. A= 13.81, B= -0.0234
154SM CL BAND(B) Octupole-vibrational band. A= 10.58, B= -0.0413
154SM XR154PM B- DECAY (1.73 MIN)
154SM XS154PM B- DECAY (2.68 MIN)
154SM XT154PM(3HE,3H)
154SM XU154SM(N,N'G)
154SM XVCOULOMB EXCITATION
154SM XWINELASTIC SCATTERING
154SM XX154SM(G,G')
154SM XY152SM(T,P)
154SM XZ154EU EC DECAY
154SM L 0.0 0+ STABLE A
154SM L 81.99 2 2+ 3.02 NS 4 A
154SM2 L MOMM2= -1.87 4 $ BE2= 4.32 2 $ MOMM1= +0.70 5 $
154SM CL J From E2 Coulomb excitation from 0+ ground state.
154SM L 266.74 7 4+ 172 PS 4 A
154SM2 L MOMM1= +1.35 15 $ MOMM2= -2.2 8 $ BE4= 0.305 18 $ XREF= -(TX) $
154SM CL J From multiple Coulomb excitation and expected 0+ band
154SM2CL structure.
154SM L 543.9 5 6+ 22.7 PS 6 A
154SM2 L MOMM1= +1.90 28 $ XREF= SVWY
154SM CL J From multiple Coulomb excitation and expected 0+ band
154SM2CL structure.
154SM CL MOMM1 From 72KU10 and given as g-factor of 0.317 46. (Also given
154SM2CL by 76FU06 and 78LEZA.)
154SM L 902.8 8 8+ 5.93 PS 25 A
154SM2 L XREF= V
154SM CL J From multiple Coulomb excitation and expected 0+ band
154SM2CL structure.
154SM L 921.57 11 1- 26 FS 4 B
154SM2 L XREF= RSUWX
154SM CL J From E1 excitation by (G,G') reaction.
154SM2CL partial-width data (76ME17) from (G,G') experiment.
154SM L 1012.62 12 3- B
154SM2 L BE3= 0.10 2 $ XREF= SUVW
154SM CL J From E3 Coulomb excitation.
154SM CL BE3 Weighted average of 0.11 3 (68KE04) and 0.09 2 (68VE01).
154SM2CL Other: 66SE06.

```

APPENDIX C-2 (cont'd)

Example Output for the Adopted Levels Data Set

¹⁵⁴Sm Adopted Levels

Q(β^-) = - 717.0 21; S(n)=7967.9 10; S(p)=9086 16; Q(α) = - 1197.2 25 85Wa02.

Summary of Adopted T_{1/2} Values

E(level)	Adopted	Comments
0.0	Stable	
81.9	3.02 ns 4	Weighted average of 3.03 ns 5 (67Wo06) and 3.00 ns 6 (68Ri09). Other: 2.74 ns 24 (59Bi10).
266.7	172 ps 4	Weighted average of 173 ps 5 (72Di06) and 169 ps 10 (80Jo08).
543.9	22.7 ps 6	Weighted average of 23.3 ps 7 (72Di06) and 22.7 ps 6 (average of two values in 80Jo08).
902.8	5.93 ps 25	Weighted average of 6.2 ps 6 (72Di06), 6.0 ps 4 (77Ke06), and 5.8 ps 4 (80Jo08).
921.5	26 fs 4	Calculated from level width of 0.0176 eV 24.

Summary of J π Assignments

E(level)	Adopted	β^- decay	(n,n' γ)	Coulomb excit
0.0	0 +	0 +	0 +	0 +
81.9	2 +	2 +	2 +	2 +
266.7	4 +	4 +	4 +	4 +
543.9	6 +	6 +		6 +

The data on the level energies and J π values are primarily from the ¹⁵⁴Pm β^- decay.

Cross Reference (XREF) Flags

A ¹⁵⁴ Pm β^- Decay (1.73 min)	F Inelastic Scattering
B ¹⁵⁴ Pm β^- Decay (2.68 min)	G ¹⁵⁴ Sm(γ,γ')
C ¹⁵⁴ Pm(³ He, ³ H)	H ¹⁵² Sm(t,p)
D ¹⁵⁴ Sm(n,n' γ)	I ¹⁵⁴ Eu ϵ Decay
E Coulomb Excitation	

E(level) \ddagger	J π	XREF	T _{1/2}	Comments
0.0 \ddagger	0 +		stable	
81.99 \ddagger 2	2 +		3.02 ns 4	Q = - 187 4; B(E2) = 4.32 2; μ = + 0.70 5 J π : From E2 Coulomb excitation from 0+ ground state.
266.74 \ddagger 7	4 +	C G	172 ps 4	μ = +1.35 15; Q = - 2.2 8; B(E4)=0.305 18. J π : from multiple Coulomb excitation and expected 0+ band structure.
543.9 \ddagger 5	6 +	B EF H	22.7 ps 6	μ = +1.90 28. J π : From multiple Coulomb excitation and expected 0+ band structure. μ : From 72Ku10 and given as a g--factor of 0.317 46 (Also given by 76Fu06 and 78LeZA).
902.8 \ddagger 8	8 +	E	5.93 ps 25	J π : From multiple Coulomb excitation and expected 0+ band structure.
921.57 \S 11	1 -	AB D FG	26 fs 4	J π : From E1 excitation by (γ,γ') reaction; partial-width data (76Me17) from (γ,γ') experiment.
1012.62 \S 12	3 -	B DEF		B(E3)=0.10 2. J π : From E3 Coulomb excitation. B(E3): Weighted average of 0.11 3 (68Ke04) and 0.09 2 (68Ve01). Other: 66Se06.

\ddagger From least-squares fit to γ energies.

\ddagger K π = 0 + ground-state band. A=13.81, B= - 0.0234.

\S Octupole-vibrational band. A=10.58, B= - 0.0413.

APPENDIX C-3

Example of a Decay Data Set

```

154SM 154PM B- DECAY (1.7 M) 74YA07,72TA13,71DA28 87NDS 870101
154PM P (0,1) 1.7 M 2 4.0E+3 1
154SM N 0.194 30 1.0
154SM CN NR I(G+CE)=100 TO G.S. IF IT-DECAY OF PARENT=0
154SM C OTHERS: 68DEZZ,72H008,73PR05,74BU09.
154SM CG E,RI$FROM 74YA07 UNLESS OTHERWISE STATED.
154SM CG RI(B)$DATA NOT GIVEN; DEDUCED FROM G-BRANCHING RATIOS VIA 2.7-MIN
154SM CB MEASURED BETAS: 71DA28,74YA07. OTHERS: 73PR05,72TA13,58WI42
154SM G 414.8 2 5.9 9
154SM L 0.0 0+ STABLE
154SM L 81.98 10 2+
154SM G 81.98 10 65 10 E2 4.93 C
154SMS G KC= 2.021 $LC= 2.250 $MC= 0.519 $NC+= 0.1402 $
154SM L 266.7 2 4+
154SM G 184.76 10 25 5 E2 0.27 C
154SMS G KC= 0.1925 $LC= 0.0633 $MC=0.01428 $NC+=0.00389 $
154SM CG E FROM 71DA28. OTHER: 184.6 1 (74YA07)
154SM L 921.6 2 1-
154SM B 12 2 6.5 C
154SM2 B EAV= 1250 50$
154SM CB EB=3.0 2 MEV (74YA07,71DA28) SCIN, F-K
154SM G 839.6 2 66.3 38 C
154SM G 921.6 2 44.3 27 C
154SM L 1099.7 3 0+
154SM B 5 1 6.8
154SM2 B EAV= 1170 50$
154SM G 1017.6 2 53.1 32 C
154SM L 1178.2 3 2+
154SM B 3.5 12 6 9
154SM2 B EAV= 1140 50$
154SM G 911.1 3 24.1 25 C
154SM G 1096.2 3 29.9 22 C
154SM G 1177.8 8 19.0 19 C
154SM L 1440.4 4 2+
154SM G 1173.7 100.08 5 B
154SM G 1358.6 2 1.3 4 B
154SM G 1440.3 2 1.1 4 B
154SM L 1476.1 3 (1-)
154SM B 11 2 6.2 C
154SM2 B EAV= 1000 50$
154SM G 1394.0 2 65 10 C
154SM L 1890.9 3
154SM B 3.0 5 6.5
154SM2 B EAV= 820 50$
154SM G 1808.8 3 8.4 13
154SM G 1891.0 4 7.0 14
154SM L 2089.8 3
154SM B 25 2 5.4 C
154SM2 B EAV= 740 50$
154SM G 891.5 2 35.6 23
154SM G 970.0 2 26.9 20
154SM G 1148.1 2 48.2 30 C
154SM G 1988.0 3 6.7 10
154SM G 2070.2 3 9.6 10
154SM L 2140.9 2
154SM B 36 3 5.2 C
154SM2 B EAV= 700 50$
154SM G 664.6 2 7.2 14
154SM G 700.4 6 2.5 8
154SM G 962.5 2 19.1 16 C
154SM G 1218.0 10

```


APPENDIX C-3 (cont'd)

Example for Radiation Tables

β^- radiations from ^{154}Pm β^- Decay (1.7 min) 74Ya07,72Ta13,71Da28

Others: 68DeZZ,72Ho08,73Pr05,74Bu09.

Measured β^- 's: 71Da28,74Ya07. Others: 73Pr05,72Ta13,58Wi42.

$E\beta^-$	E(level)	$I\beta^-^\dagger$	Log ft	Comments
(1380 100)	2618.2	1.0 3	6.2	av $E\beta=500$ 50.
(1410 100)	2592.3	1.8 2	8.0	av $E\beta=510$ 50.
(1650 100)	2347.7	1.8 2	6.3	av $E\beta=610$ 50.
(1860 100)	2140.9	36. 3	5.2	av $E\beta=700$ 50.
(1930 100)	2069.8	25. 2	5.4	av $E\beta=740$ 50.
(2110 100)	1890.9	3.0 5	6.5	av $E\beta=820$ 50.
(2520 100)	1476. 1	11. 2	6.2	av $E\beta=1000$ 50.
(2820 100)	1178.2	3.5 12	6.9	av $E\beta=1140$ 50.
(2900 100)	1099.7	5. 1	6.8	av $E\beta=1170$ 50.
(3080 100)	921. 6	12. 2	6.5	av $E\beta=1250$ 50.

$E\beta=3.0$ 2 MeV (74Ya07,71Da28) scin, F-K.

† For β^- intensity per 100 decays, multiply by 1.00.

$\gamma(^{154}\text{Sm})$ from ^{154}Pm β^- Decay (1.7 min) 74Ya07,72Ta13,71Da28

I_γ normalization: $I(\gamma+ce)=100$ to g.s. if IT-decay of parent=0.

$E\gamma^\ddagger$	E(level)	$I_\gamma^{\ddagger\pm}$	Mult.	α	Comments
81.9810	81.98	65 10	E2	4.93	$\alpha(K)=2.021$; $\alpha(L)=2.250$; $\alpha(M)=0.519$; $\alpha(N+)=0.1402$.
184.7610	266.7	25 5	E2	0.27	$\alpha(K)=0.1925$; $\alpha(L)=0.0633$; $\alpha(M)=0.01428$; $\alpha(N+..)=0.00389$. $E\gamma$: from 71DaZ8. Other: 184.6 1 (74Ya07).
*414.8 2		5.9 9			
664.6 2	2140.9	7.2 14			
700.4 6	2140.9	2.5 8			
839.6 2	921.6	66.3 38			
891.5 2	2069.8	35.6 23			
911.1 3	1178.2	24.1 25			
921.6 2	921.6	44.3 27			
962.5 2	2140.9	19.1 16			
970.0 2	2069.8	26.9 20			
1017.6 2	1099.7	53.1 32			
1096.2 3	1178.2	29.9 22			
1146.1 2	2069.8	48.2 30			
1173.7 10	1440.4	0.08 [§] 5			
1177.8 8	1178.2	19.0 19			
1218.0 10	2140.9				
1358.8 2	1440.4	1.3 [§] 4			
1394.0 2	1476. 1	65 10			
1440.3 2	1440.4	1.1 [§] 4			
1808.8 3	1880.9	8.4 13			
1891.0 4	1880.9	7.0 14			
1988.0 3	2069.8	6.7 10			
2058.9 2	2140.9	100			
2070.2 3	2069.8	9.6 10			
2140.9 2	2140.9	56.7 3			
2347.7 3	2347.7	9.2 9			
2510.6 2	2592.3	7.0 8			
2535.9 3	2618.2	3.3 8			
2592.0 3	2592.3	2.2 5			
2618.8 5	2618.2	2.0 8			

‡ From 74Ya07 unless otherwise noted.

† For absolute intensity per 100 decays, multiply by 0.19 3.

§ Data not given; deduced from γ -branching ratios via 2.7-min.

* γ ray not placed in level scheme.

APPENDIX D

ENSDF TRANSLATION DICTIONARY

The publication programs translate the text of comments (**CTEXT** in Section **III.B**) from the computer-readable **ENSDF** input into printed output for *Nuclear Data Sheets*.

The translation dictionary, as of January 1987, is given below. It is given sorted by the output (translation) in (Appendix D-I) and by the ENSDF input (Appendix D-II). The dictionary is constantly enlarged and improved as new needs are encountered.

APPENDIX D-1 (Cont.)

ENSDF DICTIONARY— ordered by output

<u>Translation</u>	<u>ENSDF</u>	<u>Translation</u>	<u>ENSDF</u>
A	^A	B(M1)(W.u.)	BM1W
av E β	EAV	B(M2)	BM2
Ay	AY	B(M2)(W.u.)	BM2W
A(θ)	A(THETA)	B(M3)	BM3
A-N	A-N	B(M3)(W.u.)	BM3W
A ^{1/3}	A**1/3	B(M4)	BM4
A ^{2/3}	A**2/3	B(M4)(W.u.)	BM4W
A ₀	A0	B(M5)(W.u.)	BM5W
a ₀	a0	B \times ρ	B*RHO
A ₁	A1	C	C
A ₂₂	A22	CCBA	CCBA
A ₂	A2	ce	CE
A ₂ P ₂	A2P2	CERN	CERN
A ₂ /A ₀	A2/A0	ce(K)	CEK
A ₃	A3	ce(K)/(γ +ce)	K/T
A ₄	A4	ce(L)	CEL
B	^B	ce(L)/(γ +ce)	L/T
Be	Be	ce(L1)	CEL1
BE(L)	BE(L)	ce(L2)	CEL2
BF ₃	BF3	ce(L3)	CEL3
BG ₀	BG0	ce(M)	CEM
BJ ²	BJ**2	ce(M)/(γ +ce)	M/T
BM(L)	BM(L)	ce(M+)/(γ +ce)	M+/T
Branching	BR	ce(M1)	CEM1
branching uncertainty	DBR	ce(M2)	CEM2
B(EL)	BEL	ce(M3)	CEM3
B(EL)(W.u.)	BELW	ce(M4)	CEM4
B(E0)	B(E0)	ce(M5)	CEM5
B(E0)	BE0	ce(N)	CEN
B(E0)(W.u.)	BE0W	ce(N+)/(γ +ce)	N+/T
B(E1)	B(E1)	ce(N1)	CEN1
B(E1)	BE1	ce(N2)	CEN2
B(E1)(W.u.)	BE1W	ce(N3)	CEN3
B(E2)	B(E2)	ce(N4)	CEN4
B(E2)	BE2	ce(N5)	CEN5
B(E2)(W.u.)	BE2W	ce(0)	CE0
B(E2) \uparrow	BE2UP	ce(0)+ce(P)	CE0+CEP
B(E3)	B(E3)	ce β	CEB
B(E3)	BE3	ce γ	CEG
B(E3)(W.u.)	BE3W	Cm	Cm
B(E3) \uparrow	BE3UP	cm ²	CM2
B(E4)	B(E4)	cm ³	CM3
B(E4)	BE4	Co	CO
B(E4)(W.u.)	BE4W	configuration	CONF
B(E4) \uparrow	BE4UP	configuration=	CONF=
B(E5)	BE5	cos ² θ	COS2TH
B(E5)(W.u.)	BE5W	Coul	COUL
B(E6)	BE6	CP	CP
B(E6)(W.u.)	BE6W	CsI	CSI
B(E7)	BE7	c.m.	C.M.
B(E8)	BE8	C ² S	C2S
B(J)	B(J)	D	^D
B(ML)	BML	DC0	DC0
B(ML)(W.u.)	BMLW	DC0Q	DC0Q
B(M1)	BM1	dE/dx	DE/DX
		DPAC	DPAC

APPENDIX D-1 (Cont.)

ENSDF DICTIONARY— ordered by output

<u>Translation</u>	<u>ENSDF</u>	<u>Translation</u>	<u>ENSDF</u>
DPAD	DPAD	fm ⁻¹	FM-1
DSA	DSA	fm ²	FM**2
DSAM	DSAM	fm ⁴	FM**4
DWBA	DWBA	FWHM	FWHM
DWIA	DWIA	F-K	F-K
DWUCK	DWUCK	GDR	GDR
D+Q	D+Q	Geiger	GEIGER
D+(Q)	D+(Q)	GeV	GEV
d ³ He	D3HE	Ge(Li)	GELI
dγ	DG	GM	GM
dσ	DSIGMA	GMR	GMR
dσ/dΩ	DS/DW	GQR	GQR
dΩ	DOMEGA	gT	G*T
E	E	GTOL	GTOL
Ee	EE	gT _{1/2}	GT1/2
EL	EL	gwΓγ ₀	G*W*WIDTHG0
ENDOR	ENDOR	g-factor	G-FACTOR
ENSDF	ENSDF	g.s.	GS
EPR	EPR	gΓ	G*WIDTH
ESR	ESR	gΓ ²	G*WIDTHG0**2
eV	EV	gΓ ² _{γ₀}	*G*WIDTHG0**2
even-A	EVEN-A	gΓ ² _{γ₀}	*G*WIDTHG0**2
EWSR	EWSR	gΓγ ₀	G*WIDTHG0
e'(θ)	E'(THETA)	HF	HF
E(cc)	ECE	HI	HI
E(d)	E(D)	HPGE	HPGE
E(e)	E(E)	ħ ω	HOMEGA
E(n)	EN	h ²	H**2
E(n)	E(N)	I	I
E(p)	EP	IAR	IAR
E(p)	E(P)	IAS	IAS
E(t)	E(T)	IBA	IBA
E(α)	E(A)	IBS	IBS
E-E	E-E	Ice	ICE
E/ΔE	E/DE	IMPAC	IMPAC
E0	E0	IPAC	IPAC
E1	E1	ISOLDE	ISOLDE
E2	E2	IT decay	IT DECAY
E3	E3	I(γ + ce)	TI
E4	E4	Iα	IA
E5	E5	Iβ	IB
E6	E6	Iβ normalization	NB
E7	E7	iβ ⁺	IB+
E8	E8	Iβ ⁻	IB-
E9	E9	Iε	IE
e ⁺	E+	Iγ	IG
E ^{1/2}	E**1/2	Iγ	RI
E ²	E**2	Iγ normalization	NR
Eα	EA	IγEγ	IG*EG
Eβ	EB	J	J
EΔE	EDE	JKπ	JKP
Eε	EEC	Jmax	JMAX
Eγ	EG	Jmin	JMIN
Eγ ⁵	EG**5	JOSEF	JOSEF
fm	FM	JULIE	JULIE
fm ⁻¹	FM**1		

APPENDIX D-1 (Cont.)

ENSDF DICTIONARY— ordered by output

<u>Translation</u>	<u>ENSDF</u>	<u>Translation</u>	<u>ENSDF</u>
J _f	JF	Lα ₂ x ray	XLA2
J _i	JI	Lβ ₁ x ray	XLB1
J ₀	J0	Lβ ₂ x ray	XLB2
J ₁	J1	Lγ ₁ x ray	XLG1
J ₂	J2	Lγ ₂ x ray	XLG2
Jπ	JPI	M	M
K	K	M x ray	XM
keV	KEV	mb	MB
kG	KG	mb/sr	MB/SR
KLL	KLL	MEDLIST	MEDLIST
kOe	KOE	MeV	MEV
K-O ₂₃ x ray	XKO23	meV	MILLI-EV
K-O ₂ x ray	XKO2	mg/cm ²	MG/CM2
K-O ₃ x ray	XKO3	ms	MS
K/L+M	K/L+M	M1	M1
K x ray	XK	M2	M2
Kα ₁ x ray	XKA1	M3	M3
Kα ₂ x ray	XKA2	M4	M4
Kα x ray	XKA	M5	M5
Kβ ₁ x ray	XKB1	N	^N
Kβ ₁ ' x ray	XKB1P	NaI	NAI
Kβ ₂ x ray	XKB2	NBS	NBS
Kβ ₂ ' x ray	XKB2P	nb / sr	NB/SR
Kβ ₃ x ray	XKB3	NC ² S	NC2S
Kβ ₄ x ray	XKB4	Ne	Ne
Kβ ₅ x ray	XKB5	NMR	NMR
Kβ ₅ ^I x ray	XKB5I	Note:	NOTE:
Kβ ₅ ^{II} x ray	XKB5II	NQR	NQR
Kβ x ray	XKB	NSσ	NS*SIGMA
Kπ	KPI	NX	NX
L	L	N-Z	N-Z
LAMPF	LAMPF	nγ	NG
Larmor	LARMOR	nγγ	NGG
Li	LI	N×σ	N*SIGMA
log ft	LOGFT	O	O
log f ¹ t	LOGF1T	odd-A	ODD-A
log f ^{1u} t	LOGF1UT	OSIRIS	OSIRIS
L(n)	LN	P	^P
L(p)	LP	PAC	PAC
L1	L1	PAD	PAD
L2	L2	PWBA	PWBA
L3	L3	PWIA	PWIA
L ₁ x ray	XL1	p(θ)	P(THETA)
L ₂ x ray	XL2	p-width	P-WIDTH
L ₃ x ray	XL3	pα	PALPHA
L _l x ray	XLL	pγ	PG
Lα x ray	XLA	pγγ	PGG
Lβ x ray	XLB	Q	MOME2
Lγ x ray	XLG	Q	Q
L x ray	XL	Q(g.s.)	QP
Lα ₁ x ray	XLA1	Q(α)	QA
		Q(β ⁻)	Q-
		Q(ε)	Q+ ₋
		Q+O	Q+O
		Q3D	Q3D

APPENDIX D-1 (Cont.)

ENSDF DICTIONARY— ordered by output

<u>Translation</u>	<u>ENSDF</u>	<u>Translation</u>	<u>ENSDF</u>
R	R	$2\beta^-$	$2B^-$
RPA	RPA	^{12}C	^{12}C
RUL	RUL	^{14}C	^{14}C
r^2	R**2	α	A
r_0	R0	β^+	B+_
S	^S	β^+n	B+N
S value	S VALUE	β^+p	B+P
SF	SF	$\beta^+\alpha$	B+A
Si(Li)	SILI	β^-	B-_
Sn	Sn	β^-n	B-N
SOREQ	SOREQ	β^-p	B-P
syst	SY	ϵ	EC
S'	S'	ϵp	ECP
S(n)	SN	γ	G
S(p)	SP	(fragment) γ	FG
S(α)	SA	(H,t)	(H,T)
S-factor	S-FACTOR	(K x ray) γ	XKG
S-value	S-VALUE	(t)	(T)
s-wave	S-WAVE	(α)	(A)
s^{-1}	S-1	(α)(ce) ,	ACE
T	ISPIN	(α)(K x ray)	AXK
T	TEMP	(β)	(B)
T	^T	(θ ,H)	(THETA,H)
th	TH	(\uparrow)	(UP)
Ti	TI	(\downarrow)	(DOWN)
tof	TOF	2J	2J
TPAD	TPAD	$2\beta^-$	2B-
TRISTAN	TRISTAN	4π	4PI
TRIUMPH	TRIUMPH	$4\pi\beta$	4PIB
T20	T20	$4\pi\beta\gamma$	4PIBG
T21	T21	$4\pi\gamma$	4PIG
T22	T22	<	LT
T_z	ISPINZ	>	GT
$T_{1/2}$	T	°	DEG
$t\gamma$	TG	$\langle r^2 \rangle$	AVRSQ
U	U	$\Delta\langle r^2 \rangle$	DAVRSQ
UNISOR	UNISOR	e.g.	E.G.
V	V	i.e.	I.E.
W	W	L	**L
$w(\theta)g\Gamma_\gamma0$	W(THETA)*G*WIDTHG0	- 1	**_1
W.u..	W.U.	- 3	**_3
X	^X	- 4	**_4
XX	XX	1 / 3	**1 / 3
x-ray	X-RAY	$^{12}C\gamma$	C12G
$X\gamma$	XG	α	ALPHA
Y	YTTRIUM	α	CC
Y	^Y	α decay	A DECAY
Z	Z	α syst	A SYST
%EWSR	%EWSR	α (exp)	ECC
%E0	%E0	α (K)	KC
%E2	%E2	α (K)exp	EKC
%IT	%IT	α (L)	LC
%I β	%IB		
%I γ	%RI		
%M1	%M1		
%SF	%SF		

APPENDIX D-1 (Cont.)

ENSDF DICTIONARY— ordered by output

<u>Translation</u>	<u>ENSDF</u>	<u>Translation</u>	<u>ENSDF</u>
$\alpha(L)\text{exp}$	ELC	β_3	B3
$\alpha(L+\dots)\text{exp}$	ELC+	β_3R	B3*R
$\alpha(L1)$	L1C	β_{42}	B42
$\alpha(L1)\text{exp}$	EL1C	β_4	B4
$\alpha(L2)$	L2C	β_{4R}	B4*R
$\alpha(L2)\text{exp}$	EL2C	β_5	B5
$\alpha(L3)$	L3C	β_6	B6
$\alpha(L3)\text{exp}$	EL3C	β_7	B7
$\alpha(M)$	MC	$\beta\alpha$	BA
$\alpha(M)\text{exp}$	EMC	$\beta\beta$	BB
$\alpha(M+\dots)$	MC+	$\beta\gamma$	BG
$\alpha(M+\dots)\text{exp}$	EMC+	$\beta\gamma_n$	BGN
$\alpha(M1)$	M1C	$\beta\gamma\gamma$	BGG
$\alpha(M1)\text{exp}$	EM1C	δ	MR
$\alpha(M2)$	M2C	ΔA	DA
$\alpha(M2)\text{exp}$	EM2C	ΔA_2	DA2
$\alpha(M3)$	M3C	ΔA_4	DA4
$\alpha(M3)\text{exp}$	EM3C	ΔE	DE
$\alpha(M4)$	M4C	$\Delta I(\gamma+ce)$	DTI
$\alpha(M5)$	M5C	$\Delta I\alpha$	DIA
$\alpha(N)$	NC	$\Delta I\beta$	DIB
$\alpha(N)\text{exp}$	ENC	$\Delta I\epsilon$	DIE
$\alpha(N+\dots)$	NC+	$\Delta I\gamma$	DRI
$\alpha(N+\dots)\text{exp}$	ENC+	$\Delta I\gamma(\%)$	PRI
$\alpha(N1)$	N1C	ΔJ	DJ
$\alpha(N2)$	N2C	$\Delta J\pi$	DJPI
$\alpha(N2)\text{exp}$	EN2C	ΔK	DK
$\alpha(N3)$	N3C	ΔL	DL
$\alpha(N3)\text{exp}$	EN3C	$\Delta Q(\alpha)$	DQA
$\alpha\text{-syst}$	A-SYST	$\Delta Q(\epsilon)$	DQ+
$\alpha\alpha$	AA	ΔS	DS
$\alpha\gamma$	AG	ΔS_n	DSN
β	B	$\Delta S(p)$	DSP
β	BETA	ΔT	DISPIN
β_c	BC	$\Delta T_{1/2}$	DT
β_{ce}	BCE	$\Delta(HF)$	DHF
βe^-	BE-	$\Delta(\log ft)$	DFT
β_n	BN	$\Delta(\beta\text{-normalization})$	DNB
β_p	BP	$\Delta(\gamma\text{-normalization})$	DNR
βR	B*R	Δ	DELTA
$\beta\text{-vibrational}$	B-VIBRATIONAL	δ^2	MR**2
β^+	B+	$\Delta\alpha$	DCC
β^-	B-	$\Delta\delta$	DMR
β_L	BL	$\Delta\pi$	DPI
β_{LR}	BL*R	ϵ	EC
$\beta_{LRA}^{1/3}$	BL*R*A**(1/3)	ϵ	EPSILON
β_0	B0	ϵB	EPSILONB
β_{12}	B12	$\epsilon B(E2)\uparrow$	EBE2UP
β_1	B1	$\epsilon B(E3)\uparrow$	EBE3UP
β_{20}	B20	ϵK	CK
β_{22}	B22	$\epsilon K(\text{exp})$	ECK
β_2	B2	ϵL	CL
β_{2R}	B2*R	$\epsilon L(\text{exp})$	ECL
β_{30}	B30	ϵM	CM
		ϵN	CN

APPENDIX D-1 (Cont.)

ENSDF DICTIONARY— ordered by output

<u>Translation</u>	<u>ENSDF</u>	<u>Translation</u>	<u>ENSDF</u>
Φ	PHI	μb	UB
γ	G	$\mu\text{b/s r}$	UB/SR
γ	GAMMA	$\mu\text{b}\times\text{MeV}$	UB*MEV
Γ	WIDTH	μg	UG
γ_{ce}	GCE	$\mu\text{g/cm}$	UG/CM
$\gamma_{\text{e-}}$	GE-	μs	US
γ_{n}	GN	μ^-	MU-
$\gamma_{\text{p}'}$	GP'	ν	NU
γ_{X}	GX	π	PI
γ/α	G/A	π^-	PI-
Γ^2	WIDTH**2	$\pi\beta$	PIB
$\Gamma_{\gamma 0}^2$	WIDTHG0**2	$\pi\beta\gamma$	PIBG
γ^{\pm}	G+-	$\pi\gamma$	PIG
$\Gamma_{\text{n}0}$	WIDTHNO	θ	THETA
Γ_{n}	WIDTHN	θ_1	THETA1
$\Gamma_{\text{p}'}$	WIDTHP'	θ_2	THETA2
$\Gamma_{\text{p}0}$	WIDTHP0	$\theta\gamma$	THETAG
$\Gamma_{\text{p}1}$	WIDTHP1	ρ	RHO
$\Gamma_{\text{p}2}$	WIDTHP2	ρ^2	RHO**2
Γ_{p}	WIDTHP	σ	SIGMA
Γ_{p}	*WIDTHP	Σ	SUMOF
$\Gamma_{\gamma 0}$	WIDTHG0	σ_{n}	SIGMAN
Γ_{γ}	WIDTHG	σ_0	SIGMA(0)
Γ_{α}	WIDTHA	σ_{γ}	SIGMAG
$\gamma\beta$	GB	σ_{ν}	SIGMANU
$\gamma\gamma$	GG	$\sigma\times\Delta E$	SIGMA*DE
$\gamma\gamma\tau$	GGG	τ	TAU
χ	GGT	ω	OMEGA
χ^2	CHI	$\omega^2\tau$	OMEGA**2*TAU
$\epsilon\text{M}(\text{exp})$	CHI**2	$\omega\tau$	OMEGA*T
$\epsilon\text{N}(\text{exp})$	ECM	ψ	PSI
κ	ECN	\times	*
λ	KAPPA	\leq	LE
μ	LAMBDA	\neq	NE
μ	MOMM1	\geq	GE
μ	MU	\approx	AP
		∞	INFNT

ENSDF

11/2(505)
 CONF=(N,NLJ)
 CONF=((P,7/2(633))(P,3/2(521))(N,3/2(621))
 CONF=(N,NLJ,-1)
 CONF=(N,1G9/2)
 CONF=(N,3G9/2,+3,23/2-)
 CONF=(N,3P1/2,-1)
 CONF=((208Pb 3-)(P,1H9/2))15/2+
 CONF=(P,1G9/2)
 CONF=((P,1H9/2,+2,8+)(N,2F5/2,-3,11/2-))25/2-
 CONF=(P,3G9/2,+3,23/2-)

TRANSLATION

11/2[505]
 configuration=(v nlj)
 configuration=((π 7/2[633])(π 3/2[521])(v 3/2[621]))
 configuration=(v nlj)⁻¹
 confieuration=(v 1g9/2)
 configuration=(v 3g9/2)⁺³23/2-
 configuration=(v 3p1/2)⁻¹
 configuration=((²⁰⁸Pb 3-)(π 1h9/2))15/2+
 configuration=(π 1g9/2)
 configuration=((π 1h9/2)²8+(v 2f5/2)⁻³11/2-)25/2-
 configuration=(π 3g9/2)⁺³23/2-

APPENDIX D-2 (cont.)

ENSDF DICTIONARY—ordered by input

<u>ENSDF</u>	<u>Translation</u>	<u>ENSDF</u>	<u>Translation</u>
A DECAY	α decay	BG	$\beta\gamma$
A SYST	α syst	BGG	$\beta\gamma\gamma$
AA	$\alpha\alpha$	BGN	$\beta\gamma n$
ACE	$(\alpha)(ce)$	BGO	BGO
AG	$\alpha\gamma$	BJ**2	BJ^2
ALPHA	α	BL	β_L
AP	\approx	BL*R	$\beta_L R$
AVRSQ	$\langle r^2 \rangle$	BL*R*A**(1/3)	$\beta_L R A^{1/3}$
AXK	$(\alpha)(K \text{ x ray})$	BML	B(ML)
AY	Ay	BMLW	B(ML)(W.u.)
A(THETA)	A(θ)	BM(L)	BM(L)
A**1/3	$A^{1/3}$	BM1	B(M1)
A**2/3	$A^{2/3}$	BM1W	B(M1)(W.u.)
A-N	A-N	BM2	B(M2)
A-SYST	α -syst	BM2W	B(M2)(W.u.)
A0	A_0	BM3	B(M3)
a 0	a 0	BM3W	B(M3)(W.u.)
A1	A_1	BM4	B(M4)
A2	A_2	BM4W	B(M4)(W.u.)
A2P2	$A_2 P_2$	BM5W	B(M5)(W.u.)
A2/A0	A_2/A_0	BN	βn
A22	A_{22}	BP	βp
A3	A_3	BR	Branching
A4	A_4	B(E0)	B(E0)
B	β	B(E1)	B(E1)
BA	$\beta\alpha$	B(E2)	B(E2)
BB	$\beta\beta$	B(E3)	B(E3)
BC	βc	B(E4)	B(E4)
BCE	βce	B(J)	B(J)
Be	Be	B*R	BR
BEL	B(EL)	B*RHO	$B \times \rho$
BELW	B(EL)(W.u.)	B+	β^+
BETA	β	B-VIBRATIONAL	β -vibrational
BE(L)	BE(L)	B-	β^-
BE-	βe^-	B0	β_0
BE0	B(E0)	B1	β_1
BE0W	B(E0)(W.u.)	B12	β_{12}
BE1	B(E1)	B2	β_2
BE1W	B(E1)(W.u.)	B2*R	$\beta_2 R$
BE2	B(E2)	B20	β_{20}
BE2UP	B(E2) \uparrow	B22	β_{22}
BE2W	B(E2)(W.u.)	B3	β_3
BE3	B(E3)	B3*R	$\beta_3 R$
BE3UP	B(E3) \uparrow	B30	β_{30}
BE3W	B(E3)(W.u.)	B4	β_4
BE4	B(E4)	B4*R	$\beta_4 R$
BE4UP	B(E4) \uparrow	B42	β_{42}
BE4W	B(E4)(W.u.)	B5	β_5
BE5	B(E5)	B6	β_6
BE5W	B(E5)(W.u.)	B7	β_7
BE6	B(E6)	C	C
BE6W	B(E6)(W.u.)	CC	α
BE7	B(E7)	CCBA	CCBA
BE8	B(E8)		
BF3	BF3		

ENSDF DICTIONARY—ordered by input

<u>ENSDF</u>	<u>Translation</u>	<u>ENSDF</u>	<u>Translation</u>
CEB	ce β	DHF	$\Delta(\text{HF})$
CEG	ce γ	DIA	$\Delta I\alpha$
CEK	ce(K)	DIB	$\Delta I\beta$
CEL	ce(L)	DIE	$\Delta I\epsilon$
CEL1	ce(L1)	DISPIN	ΔT
CEL2	ce(L2)	DJ	ΔJ
CE	ce	DJPI	$\Delta J\pi$
CEL3	ce(L3)	DK	ΔK
CEM	ce(M)	DL	ΔL
CEM1	ce(M1)	DMR	$\Delta\delta$
CEN2	ce(M2)	DNB	$\Delta(\beta\text{-normalization})$
CEM3	ce(M3)	DNR	$\Delta(\gamma\text{-normalization})$
CEM4	ce(M4)	DOMEGA	$d\Omega$
CEM5	ce(M5)	DPAC	DPAC
CEN	ce(N)	DPAD	DPAD
CEN1	ce(N1)	DPI	$\Delta\pi$
CEN2	ce(N2)	DQA	$\Delta Q(\alpha)$
CEN3	ce(N3)	DQ+	$\Delta Q(\epsilon)$
CEN4	ce(N4)	DRI	$\Delta I\gamma$
CEN5	ce(N5)	DS	ΔS
CE0	ce(0)	DSA	DSA
CE0+CEP	ce(0)+ce(P)	DSAM	DSAM
CERN	CERN	DSIGMA	$d\sigma$
CHI	χ	DSN	ΔS_n
CHI**2	χ^2	DSP	$\Delta S(p)$
CK	ϵK	DS/DW	$d\sigma/d\Omega$
CL	ϵL	DT	$\Delta T_{1/2}$
Cm	Cm	DTI	$\Delta I(\gamma+ce)$
CM	ϵM	DWBA	DWBA
CM2	cm^2	DWIA	DWIA
CM3	cm^3	DWUCK	DWUCK
CN	ϵN	D+Q	D+Q
CO	Co	D+(Q)	D+(Q)
CONF	configuration	D3HE	$d^3\text{He}$
CONF=	configuration=	E	E
COS2TH	$\cos^2\theta$	EA	$E\alpha$
COUL	Coul	EAV	$av E\beta$
CP	CP	EB	$E\beta$
CSI	CsI	EBE2UP	$\epsilon B(E2)\uparrow$
C.M.	c.m.	EBE3UP	$\epsilon B(E3)\uparrow$
C12G	$^{12}\text{C}\gamma$	EC	ϵ
C2S	C^2S	ECC	$\alpha(\text{exp})$
DA	ΔA	ECE	$E(\text{ce})$
DAVRSQ	$\Delta\langle r^2 \rangle$	ECK	$\epsilon K(\text{exp})$
DA2	ΔA_2	ECL	$\epsilon L(\text{exp})$
DA4	ΔA_4	ECU	$\epsilon M(\text{exp})$
DBR	branching uncertainty	ECN	$\epsilon N(\text{exp})$
DCC	$\Delta\alpha$	EDE	$E\Delta E$
DCO	DCO	EE	Ee
DCOQ	DCOQ	EEC	Ee
DE	ΔE	EG	$E\gamma$
DEG	$^\circ$	EG**5	$E\gamma^5$
DELTA	Δ	EKC	$\alpha(K)\text{exp}$
DE/DX	dE/dx	EL	EL
DFT	$\Delta(\log ft)$	ELC	$\alpha(L)\text{exp}$
DG	$d\gamma$	ELC+	$\alpha(L+\dots)\text{exp}$
		BLIC	$\alpha(L1)\text{exp}$

APPENDIX D-2 (cont.)

ENSDF DICTIONARY—ordered by input

<u>ENSDF</u>	<u>Translation</u>	<u>ENSDF</u>	<u>Translation</u>
EL2C	$\alpha(L2)exp$	GCE	γ_{ce}
EL3C	$\alpha(L3)exp$	GDR	GDR
EMC	$\alpha(M)exp$	GE	\geq
EMC+	$\alpha(M+...)exp$	GEIGER	Geiger
EM1C	$\alpha(M1)exp$	GELI	Ge(Li)
EM2C	$\alpha(M2)exp$	GEV	GeV
EM3C	$\alpha(M3)exp$	GE-	γe^-
EN	E(n)	GG	$\gamma\gamma$
ENC	$\alpha(N)exp$	GGG	$\gamma\gamma\gamma$
ENC+	$\alpha(N+...)exp$	GGT	$\gamma\gamma\tau$
ENDOR	ENDOR	GM	GM
ENSDF	ENSDF	GMR	GMR
EN2C	$\alpha(N2)exp$	GN	γn
EN3C	$\alpha(N3)exp$	CP'	$\gamma p'$
EP	E(p)	GQR	GQR
EPR	EPR	GS	g.s.
EPSILON	ϵ	GT	$>$
EPSILONB	ϵB	GTOL	GTOL
ESR	ESR	GT1/2	$gT_{1/2}$
EV	eV	GX	γX
EVEN-A	even-A	G*T	gT
EWSR	EWSR	G*WIDTH	$g\Gamma$
E'(THETA)	$e'(\theta)$	G*WIDTHG0	$g\Gamma\gamma_0$
E(A)	E(α)	G*WIDTHG0**2	$g\Gamma\gamma_0^2$
E(D)	E(d)	G*W*WIDTHG0	$gw\Gamma\gamma_0$
E(E)	E(e)	G+-	γ^\pm
E(N)	E(n)	G-FACTOR	g-factor
E(P)	E(p)	G/A	γ/α
E(T)	E(t)	HF	HF
E**1/2	$E^{1/2}$	HI	HI
E**2	E^2	HOMEGA	$\hbar\omega$
E+	e^+	HPGE	HPGE
E-E	E-E	H**2	h^2
E.G.	e.g.	I	I
E/DE	$E/\Delta E$	IA	$I\alpha$
E0	E0	IAR	IAR
E1	E1	IAS	IAS
E2	E2	IB	$I\beta$
E3	E3	IBA	IBA
E4	E4	IBS	IBS
E5	E5	IB+	$I\beta^+$
E6	E6	IB-	$I\beta^-$
E7	E7	ICE	Ice
E8	E8	IE	$I\epsilon$
E9	E9	IG	$I\gamma$
FG	(fragment) γ	IG*EG	$I\gamma E\gamma$
FM	fm	IMPAC	IMPAC
FM**-1	fm^{-1}	INFNT	∞
FM**2	fm^2	IPAC	IPAC
FM**4	fm^4	ISOLDE	ISOLDE
FM-1	fm^{-1}	ISPIN	T
FWHM	FWHM	ISPINZ	T_z
F-K	F-K	IT DECAY	IT decay
G	γ	I.E.	i.e.
GAMMA	γ	J	J
GB	$\gamma\beta$		

APPENDIX D-2 (cont.)

ENSDF DICTIONARY—ordered by input

<u>ENSDF</u>	<u>Translation</u>	<u>ENSDF</u>	<u>Translation</u>
JF	J _f	MU-	μ ⁻
JI	J _i	M+/T	ce(M+)/((γ+ce)
JKP	JKπ	M/T	ce(M)/((γ+ce)
JMAX	Jmax	M1	M1
JMIN	Jmin	M1C	α(M1)
JOSEF	JOSEF	M2	M2
JPI	Jπ	M2C	α(M2)
JULIE	JULIE	M3	M3
J0	J 0	M3C	α(M3)
J1	J 1	M4	M4
J2	J 2	M4C	α(M4)
K	K	M5	M5
KAPPA	κ	M5C	α(M5)
KC	α(K)	NAI	NaI
KEV	keV	NB	Iβ normalization
KG	kG	NBS	NBS
KLL	KLL	NB/SR	nb/sr
KOE	kOe	NC	α(N)
KPI	Kπ	NC+	α(N+...)
K/L+M	K/L+M	NC2S	NC ² S
K/T	ce(K)/((γ+ce)	Ne	Ne
L	L	NE	≠
LAMBDA	λ	NC	nγ
LAMPF	LAMPF	NCG	nγγ
LARMOR	Larmor	NMR	NMR
LC	α(L)	NOTE	Note
LE	≤	NQR	NQR
LI	Li	NR	Iγ normalization
LN	L(n)	NS-SIGMA	NSσ
LOGFT	log ft	NU	v
LOGF1T	log f ¹ t	NX	NX
LOGF1UT	log f ¹ U _t	N*SIGMA	N×σ
LP	L(p)	N+/T	ce(N+)/((γ+ce)
LT	<	N-Z	N-Z
L/T	ce(L)/((γ+ce)	N1C	α(N1)
L1	L1	N2C	α(N2)
L1C	α(L1)	N3C	α(N3)
L2	L2	O	O
L2C	α(L2)	ODD-A	odd-A
L3	L3	OMEGA	ω
L3C	α(L3)	OMEGA*T	ωτ
M	M	OMEGA**2*TAU	ω ² τ
MB	mb	OSIRIS	OSIRIS
MB/SR	mb/sr	PAC	PAC
MC	α(M)	PAD	PAD
MC+	α(M+...)	PALPHA	pα
MEDLIST	MEDLIST	PG	pγ
MEV	MeV	PGG	pγγ
MG/CM2	mg/cm ²	PHI	φ
MILLI-EV	meV	PI	π
MOME2	Q	PIB	πβ
MOMM1	μ	PIBG	πβγ
MR	δ	PIG	πγ
MR**2	δ ²	PI-	π ⁻
MS	ms	PRI	ΔIγ(%)
MU	μ	PSI	ψ
		PWBA	PWBA

APPENDIX D-2 (cont.)

ENSDF DICTIONARY—ordered by input

<u>ENSDF</u>	<u>Translation</u>	<u>ENSDF</u>	<u>Translation</u>
PWIA	PWIA	T22	T22
P(THETA)	p(θ)	U	U
P-WIDTH	p-width	UB	μb
Q	Q	UB*MEV	$\mu\text{b}\times\text{MeV}$
QA	Q(α)	UB/SR	$\mu\text{b}/\text{sr}$
QP	Q(g.s.)	UC	μg
Q+O	Q+O	UG/CM	$\mu\text{g}/\text{cm}$
Q+_	Q(ϵ)	UNISOR	UNISOR
Q-	Q(β^-)	US	μs
Q3D	Q3D	V	V
R	R	W	W
RHO	ρ	WIDTH	Γ
RHO**2	ρ^2	WIDTHA	Γ_α
RI	I γ	WIDTHG	Γ_γ
RPA	RPA	WIDTHG0	$\Gamma_{\gamma 0}$
RUL	RUL	WIDTHG0**2	$\Gamma_{\gamma 0}^2$
R**2	r ²	WIDTHN	Γ_n
R0	r ₀	WIDTHN0	Γ_{n0}
S VALUE	S value	WIDTHP	Γ_p
SA	S(α)	WIDTHP'	$\Gamma_{p'}$
SF	SF	WIDTHP0	Γ_{p0}
SIGMA	σ	WIDTHP1	Γ_{p1}
SIGMAG	σ_γ	WIDTHP2	Γ_{p2}
SIGMAN	σ_n	WIDTH**2	Γ^2
SIGMANU	σ_v	W(THETA)*G*WIDTHG0	w(θ)g $\Gamma_{\gamma 0}$
SIGMA(0)	σ_0	W.U.	W.u.
SIGMA*DE	$\sigma\times\Delta E$	XG	X γ
SILI	Si(Li)	XK	K x ray
Sn	Sn	XKA	K α x ray
SN	S(n)	XKA1	K α_1 x ray
SOREQ	SOREQ	XKA2	K α_2 x ray
SP	S(p)	XKB	K β x ray
SUMOF	Σ	XKB1	K β_1 x ray
SY	syst	XKB1P	K β_1' x ray
S'	S'	XKB2	K β_2 x ray
S-FACTOR	S-factor	XKB2P	K β_2' x ray
S-VALUE	S-value	XKB3	K β_3 x ray
S-WAVE	s-wave	XKB4	K β_4 x ray
S-1	s ⁻¹	XKB5	K β_5 x ray
T	T _{1/2}	XKB5I	K β_5^I x ray
TAU	τ	XKB5II	K β_5^{II} x ray
TEMP	T	XKG	(K x ray) γ
TG	t γ	XKO2	K-O ₂ x ray
TH	th	XKO23	K-O ₂₃ x ray
THETA	θ	XKO3	K-O ₃ x ray
THETAG	$\theta\gamma$	XL	L x ray
THETA1	θ_1	XLA	L α x ray
THETA2	θ_2	XLA1	L α_1 x ray
TI	I($\gamma+ce$)	XLA2	L α_2 x ray
Ti	Ti	XLB	L β x ray
TOF	tof		
TPAD	TPAD		
TRISTAN	TRISTAN		
TRIUMPH	TRIUMPH		
T20	T20		
T21	T21		

APPENDIX D-2 (cont.)

ENSDF DICTIONARY—ordered by input

<u>ENSDF</u>	<u>Translation</u>	<u>ENSDF</u>	<u>Translation</u>
XLB1	L β_1 x ray	%14C	% ¹⁴ C
XLB2	L β_2 x ray	%2B-	%2 β^-
XLG	L γ x ray	(A)	(α)
XLG1	L γ_1 x ray	(B)	(β)
XLG2	L γ_2 x ray	(DOWN)	(\downarrow)
XLL	L l x ray	(H,T)	(H,t)
XL1	L $_1$ x ray	(THETA,H)	(θ ,H)
XL2	L $_2$ x ray	(T)	(t)
XL3	L $_3$ x ray	(UP)	(\uparrow)
XM	M x ray	*	\times
XX	XX	*G*WIDTHG0**2	$g\Gamma_{\gamma 0}^2$
X-RAY	x-ray	*WIDTHP	Γ_p
YTTRIUM	Y	**L	L
Z	Z	**−1	−1
%A	% α	**−3	−3
%B+A	% $\beta^+\alpha$	**−4	−4
%B+N	% β^+n	**1/3	1/3
%B+P	% β^+p	2B−	2 β^-
%B+_	% β^+	2J	2J
%B-N	% β^-n	4PI	4 π
%B-P	% β^-p	4PIB	4 $\pi\beta$
%B−	% β^-	4PIBG	4 $\pi\beta\gamma$
%EC	% ϵ	4PIG	4 $\pi\gamma$
%ECP	% ϵp	^A	A
%EWSR	%EWSR	^B	B
%E0	%E0	^D	D
%E2	%E2	^N	N
%G	% γ	^P	P
%IB	%I β	^S	S
%IT	%IT	^T	T
%M1	%M1	^X	X
%RI	%I γ	^Y	Y
%SF	%SF		
%12C	% ¹² C		

ENSDF

11/2(505)
 CONF=(N,NLJ)
 CONF=((P,7/2(633))(P,3/2(521))(N,3/2(621))
 CONF=(N,NLJ,−1)
 CONF=(N,1G9/2)
 CONF=(N,3G9/2,+3,23/2−)
 CONF=(N,3P1/2,−1)
 CONF=((208PB 3−)(P,1H9/2))15/2+
 CONF=(P,1G9/2)
 CONF=((P,1H9/2,+2,8+)(N,2F5/2,−3,11/2−))25/2−
 CONF=(P,3G9/2,+3,23/2−)

TRANSLATION

11/2[505]
 configuration=(v nlj)
 configuration=((π 7/2[633])(π 3/2[521])(v 3/2[621]))
 configuration=(v nlj)^{−1}
 configuration=(v 1g9/2)
 configuration=(v 3g9/2)⁺³23/2−
 configuration=(v 3p1/2)^{−1}
 configuration=((²⁰⁸Pb 3−)(π 1h9/2))15/2+
 configuration=(π 1g9/2)
 configuration=((π 1h9/2²8+(v 2f5/2)^{−3}11/2−))25/2−
 configuration=(π 3g9/2)⁺³23/2−

APPENDIX E

DATA EVALUATION CENTERS

- | | |
|--|---|
| <p>a. National Nuclear Data Center
Brookhaven National Laboratory
Upton, NY 11973, U.S.A.
Contact: J. K. Tuli</p> <p>b. Nuclear Data Project
Oak Ridge National Laboratory
Oak Ridge, TN 37831, U.S.A.
Contact: M. J. Martin</p> <p>c. Isotopes Project
Lawrence Berkeley Laboratory
Berkeley, CA 94720, U.S.A.
Contact: E. Browne/R. B. Firestone</p> <p>d. Idaho National Engineering Laboratory
E.G. and G. Idaho, Inc.
P.O. Box 1625
Idaho Falls, ID 83415, U.S.A.
Contact: C. W. Reich</p> <p>e. Physics Department
University of Pennsylvania
Philadelphia, PA 19174, U.S.A.
Contact: F. Ajzenbere-Selove</p> <p>f. Center for Nuclear Structure
and Reaction Data of the U.S.S.R.
State Committee on the Utilization
of Atomic Energy U.S.S.R.
46 Ulitsa Kurchatova
Moscow, D-182, U.S.S.R.
Contact: P. E. Chukreev</p> <p>g. Data Centre
Leningrad Nuclear Physics Institute
Gatchina, Leningrad Region
188350, U.S.S.R.
Contact: I. A. Kondurov</p> <p>h. Fysisch Laboratorium
Princetonplein 5, Postbus 80.000
3508 TA Utrecht, The Netherlands
Contact: C. van der Leun</p> | <p>i. Fachinformationszentrum Energie,
Physik, Mathematik GmbH
Kernforschungszentrum
D-7514 Eggenstein-Leopoldshafen 2
F.R.G.
Contact: H. Behrens</p> <p>j. Centre d'Études Nucléaires
DRP-CPN
Cedex No. 85
F-38041 Grenoble Cedex, France
Contact: J. Blachot</p> <p>k. Division of Physics
Japan Atomic Energy Research Institute
Tokai-Mura, Naka-Gun
Ibaraki-Ken 319-11, Japan
Contact: T. Tamura</p> <p>l. Institute of Physics
University of Lund
Solvegatan 14
S-223 62 Lund, Sweden
Contact: P. Ekström</p> <p>m. Nuclear Data Project
Kuwait Institute for Scientific Research
P.O. Box 24885
Kuwait, Kuwait
Contact: S. Rab</p> <p>n. Laboratorium voor Kernfysica
Proeftuinstraat 86
B-9000 Gent, Belgium
Contact: D. De Frenne</p> <p>o. Tandem Accelerator Laboratory
McMaster University
Hamilton, Ontario L8S 4K1
Canada
Contact: J. A. Kuehner</p> <p>p. Institute of Atomic Energy
P.O. Box 275 (41), Beijing,
People's Republic of China
Contact: C. Dunjiu</p> |
|--|---|

EVALUATION RESPONSIBILITY

<u>A-Range</u>	<u>Center</u>	<u>A-Range</u>	<u>Center</u>	<u>A-Range</u>	<u>Center</u>	<u>A-Range</u>	<u>Center</u>
1-4	f	65-73	a	136-148	a	195-198	p
5-20	e	74-80	m	149,151	o	199-237	b
21-44	h	81-100	i	150,152	a	238-244(even)	f
45-50	a	101-110	j	153-162	d	239-243(odd)	b
51-56	p	111-117	n	163,165	a	245-263	b
57,58	a	118-129	k	164,166	f		
59-64	l	130-135	g	167-194	c		

APPENDIX F

Conventions, Policies and Symbols in *Nuclear Data Sheets*

- Appendix F-1 Language Style in *Nuclear Data Sheets*
- Appendix F-2 General Policies – Theory
- Appendix F-3 Bases For Spin And Parity Assignments
- Appendix F-4 Conventions
- Appendix F-5 Symbols and Abbreviations

LANGUAGE STYLE IN NUCLEAR DATA SHEETS

The *Nuclear Data Sheets* generally follow the language guidelines as contained in the *American Institute of Physics Style Manual* (AIP Publ. R-283, 1978). Some specific suggestions on punctuation, abbreviations, spelling, and grammar, as well as a discussion of some common problems follow:

a. **Abbreviations:** Use the standard abbreviations given in Appendix F-5 and the inside back cover of the journal. For example, use "excit" instead of "exc," "exp" instead of "expt," etc. Use periods with abbreviations only if given on the list. Abbreviations such as av, calc, norm, pol, rel, res, etc. should not be used in the regular text. They are intended for use in tables, equations, parenthetical expressions, reactions, etc. to conserve space.

b. **Keynumbers:** Keynumbers should be treated as plural subjects since, for example, "81Aa99 are continuing ..." is the same as "The authors of 81Aa99 are continuing..." Keynumbers in series should be in either descendine (preffered) or ascendine chronological, alphabetical, and numerical order. This should be consistent throughout an evaluation.

c. **Tense:** Past-tense verbs should be used for events which have already taken place and are not ongoing. For example "81Aa99 measured..." instead of "81Aa99 measure..." However, "81Ba12 cite..."

d. **Spelling:** When there are two or more correct spellings of a word, use the short or American spelling. For example, "analog" instead of "analogue," "behavior" instead of "behaviour," "polarization" instead of "polarisation." However, "thru," "tho," and other simplified spellings, slang, or jargon should not be used. Under l(vi) below, there is a list of some words that are always spelled solid and others that are always hyphenated. Use standard rules for hyphenating compound modifiers. Appendix B of the *AIP Style Manual* contains additional examples of the preferred spellings of words which occur frequently. *Webster's Instant Word Guide* is a very useful source for both the spelling and the hyphenation of words.

e. **Commas:** A comma precedes the conjunction before the final item in a series, unless the conjunction joins combinations. This rule would also apply to the semi-colon when used in a series. In comments listing level energy and $J\pi$, set off the $J\pi$ with commas as an appositive. For example:

1981.3, (1-,2-), is given

1993.5, 1, is given.

In a series, use semicolons after all but the last $J\pi$ (comma before conjunction). For example, 1981.3, (1-, 2-); 1992.4, 3-, 5-; and 1993.5, 1, are given.

Key numbers in comments are separated by commas immediately followed by a space. However, key numbers on formatted or continuation records or key numbers enclosed in parentheses are separated by a comma without a following blank.

f. **Person:** Smooth transitions between first and third person are acceptable.

g. **Units:** With the exception of the four standard units for energy, cross section, and electric and magnetic moments as given under the "Conventions used in Nuclear Data Sheets," appropriate units, as defined in Appendix F-5 should be given for all quantities. In eeneral, the unit should appear between the value and associated uncertainty. For example, $T_{1/2} = 13.3 \text{ min } 5$. There are a few exceptions where the unit is associated with the quantity measured or deduced. For example, $B(E2)(W.u.)=1.0 \text{ } 1$, instead of $B(E2)=1.0 \text{ W.u. } 1$.

h. **Mathematical operators and equations:** The following operators are allowed within the text: +-, =, <, >, LT, GT, LE, NE, AP. The production programs will remove the leading and trailing blanks around the operators, thus compacting an equation. The representation of complex equations often causes problems. The evaluator can code these in comments usinit the special characters as defined in Appendix A-1. If you are not sure of the proper use of special characters, write down the formula in a note to NNDC and the production staff will try to code it for you.

LANGUAGE STYLE IN NUCLEAR DATA SHEETS (cont..)

i. Miscellaneous:

- (i) Use "spin and parity" instead of "spin-parity."
- (ii) Use "uncertainty" instead of "error" unless "error" means "mistake."
- (iii) If a number contains a decimal point, there must be a number before the decimal point ("0.99" instead of ".99").
- (iv) Use the form "0.99 9" instead of "0.99±9" or "0.99±0.09" for uncertainties.
- (v) When foreign words are used, the appropriate number should be reflected in the verb and modifiers. One of the most widely used word in this category is the Latin singular word "datum," whose plural is "data." Singular "This datum is ..." would have the plural "These data are..."
- (vi) Some single words:
- | | | |
|---------------|---------------|----------------|
| antianalog | depopulation | nonstatistical |
| antiparallel | endpoint | nonuniform |
| bandhead | lifetime | nonunique |
| backbending | lineshape | overall |
| backscat-ter | nonaligned | photopeak |
| crossover | nonelastic | pickup (adj.) |
| cutoff (adj.) | nonlocal | subshell |
| deadline | nonresonant | wavelength |
| deexcitation | nonrotational | |

Some hyphenated words (Modifiers made up of two or more words are usually hyphenated)

di-pion	l-no shape	half-life
2-yes shape	half-width	96-keV level
mean-life	s-wave capture	non-negative
shell-model calculations	non-Fermi	x-ray intensity
quasi-free-electron	self-consistent	semi-infinite
up-spin		

The following are spelled as two words:

cross section
 shell model
 wave function
 x ray
 α particle
 γ ray

GENERAL POLICIES — "THEORY"

A reference "Theory 67Xy01" indicates theoretical predictions computed by the authors of 67Xy01. A reference "Theory" alone indicates a determination by the compiler of theoretical predictions described below.

Internal Conversion Coefficients

Theoretical conversion coefficients are obtained by spline interpolation (68Ha53) from tables of Hager and Seltzer (68Ha53) for the K-, L₁...3-, M₁...5-shells and of Dragoun, Plajner, and Schmutzler (71Dr11) for the (N+O+...)-shells. For the N₁...5-subshells, values are obtained by graphical interpolation from tables of Dragoun, Pauli, and Schmutzler (69Dr09). For K-, L₁...3-shells, conversion coefficients for transitions outside the E_γ, Λ-, or Z-ranges of Hager and Seltzer are obtained as follows: for E_γ ≤ 6000 keV and for Z=3, 6, 10 and 14 ≤ Z ≤ 30 interpolation from tables of Band et al. (76Ba63); for E_γ > 2600 keV, by graphical interpolation from tables of Trusov (72Tr09). For E0-transitions, K/L₁ and L₁/L₂ ratios are obtained by graphical interpolation from tables of Hager and Seltzer (69Ha61).

Angular Distribution and Correlation Coefficients

The coefficients required for analysis of directional correlation, polarization correlation, directional distribution, and polarization distribution data are obtained as described by Steffen (71St47, 71St48). In particular, we adopt the phase convention for the mixing ratio, δ, defined by the authors (70Kr03). Particle parameters required for the analysis of correlation and distribution data involving conversion electrons are obtained by graphical interpolation from tables of Hager and Seltzer (68Ha54). The expression for the deorientation coefficient required to account for intermediate unobserved mixed radiations is given by Anicin (72An20).*

A tabulation of gamma-gamma directional coefficients is given by Taylor, et al. (71Ta32). These authors use the Steffen phase convention.

Penetration Parameters

Penetration parameters required for the analysis of internal conversion data and angular correlation or distribution data involving electrons are obtained by graphical interpolation from tables of Hager and Seltzer (69Ha61).

Internal Pair Conversion Coefficients

Theoretical internal pair conversion coefficients for A=E1, M1, E2 are obtained by graphical interpolation in Z, E from tables of Lombard, et al. (68Lo16).

* As pointed out by these authors, most earlier references which discuss this coefficient define it incorrectly

β-Decay Rate Probabilities

Logft values, capture-to-positron ratios, and electron-capture ratios for allowed, first-forbidden unique, and second-forbidden unique transitions are obtained as described by Gove and Martin (71Go40). This reference also contains a tabulation of log f values and total capture-to-positron ratios for allowed and first-forbidden unique transitions.

Atomic Processes

X-ray fluorescence yields are obtained from Bambynek et al. (72Bb16) for Z ≤ 92 and from Ahmad (79Ah01) for Z > 92.

Electron binding energies for Z < 84 are taken from Bearden and Burr (67Be73) and from Porter and Freedman (78Po08) for Z > 84.

α-Decay Hindrance Factors

The α-hindrance factors (the ratio of the measured partial half-life for α-emission to the theoretical half-life) are obtained from the spin-independent equations of Preston (47Pr17). The nuclear radius for each even-even nucleus was determined by defining, for the g.s. to g.s. α-transition hindrance factor, HF = 1. For odd-A and odd-odd nuclei, the radius was chosen to be the average of the radii for the adjacent even-even nuclei (72El21). In the few cases where only one adjacent even-even radius was known, that value was corrected for the A^{1/3} mass dependence and used in the calculation.

Electromagnetic Transition Rates

The Weisskopf single-particle estimates for the half-lives of electric and magnetic multipole radiation of energy E_γ are (62B19)

$$T_{1/2W}(EL) = 0.190 \left(\frac{L}{L+1} \right) \left(\frac{3+L}{3} \right)^2 \frac{[(2L+1)!!]^2}{A^{2L/3}} \left(\frac{164.44}{E_{\gamma}(\text{MeV})} \right)^{2L+1} \times 10^{-21} \text{ s}$$

$$T_{1/2W}(ML) = 3.255 A^{2/3} T_{1/2W}(EL)$$

for nuclear radius $1.2 A^{1/3} \times 10^{-13}$ cm.

Unweighted and Weighted Averages

If $x_1 \pm \Delta x_1, x_2 \pm \Delta x_2, \dots, x_n \pm \Delta x_n$ are n independent measurements of a given quantity, Δx_i being the uncertainty in x_i, then the weighted average of these measurements is $\bar{x} \pm \Delta \bar{x}$, where

$$\bar{x} = W \sum x_i / (\Delta x_i)^2,$$

$$W = 1 / \sum (\Delta x_i)^{-2},$$

and Δx̄ is the larger of (W)^{1/2}

$$\text{and } [W \sum (\Delta x_i)^{-2} (\bar{x} - x_i)^2 / (n-1)]^{1/2}.$$

The unweighted average of these same measurements is given by $\bar{x} \pm \Delta \bar{x}$, where

$$\bar{x} = \sum x_i / n,$$

$$\Delta \bar{x} = [\sum (\bar{x} - x_i)^2 / n(n-1)]^{1/2}$$

GENERAL POLICIES—"THEORY" (cont'd)

Reference

- | | |
|--|---|
| <p>47Pr17 M. A. Preston—<i>Phys. Rev.</i> 71, 865 (1947); The Theory of Alpha Radioactivity</p> <p>52Bi97 J. M. Blatt, V. F. Weisskopf—<i>Theoretical Nuclear Physics</i>, John Wiley and Sons, Inc., New York, p. 627 (1952)</p> <p>58Ro60 M. E. Rose—<i>Internal Conversion Coefficients</i>, North-Holland Publishing Co., Amsterdam (1958)</p> <p>67Be73 J. A. Bearden, A.F. Burr—<i>Rev. Mod. Phys.</i> 39, 125 (1967); Reevaluation of X-Ray Atomic Energy Levels</p> <p>68Ha53 R. S. Hager, E. C. Seltzer—<i>Nucl. Data A4</i>, 1 (1968); Internal Conversion Tables. Part I: K-, L-, M-Shell Conversion Coefficients for Z=30 to Z=103</p> <p>68Ha54 R. S. Hager, E. C. Seltzer—<i>Nucl. Data A4</i>, 397 (1968); Internal Conversion Tables. Part II: Directional and Polarization Particle Parameters for Z=30 to Z=103</p> <p>68Lo16 R. J. Lombard, C. F. Perdriat, J. H. Brunner—<i>Nucl. Phys.</i> A110, 41 (1968); Internal Pair Formation and Multipolarity of Nuclear Transitions</p> <p>69Dr09 O. Dragoun, H. C. Pauli, F. Schmutzler—<i>Nucl. Data Tables A6</i>, 235 (1969); Tables of Internal Conversion Coefficients for N-Subshell Electrons</p> <p>69Ha61 R. S. Hager, E. C. Seltzer—<i>Nucl. Data Tables A6</i>, 1 (1969); Internal Conversion Tables. Part III: Coefficients for the Analysis of Penetration Effects in Internal Conversion and E0 Internal Conversion</p> <p>70Kr03 K. S. Krane, R. M. Steffen—<i>Phys. Rev.</i> C2, 724 (1970); Determination of the E2/M1 Multipole Mixing Ratios of the Gamma Transitions in Cd¹¹⁰</p> <p>71Dr11 O. Dragoun, Z. Plajner, F. Schmutzler - <i>Nucl. Data Tables A9</i>, 119 (1971); Contribution of Outer Atomic Shells to Total Internal Conversion Coefficients.</p> <p>71Go40 B. Gove, M. J. Martin—<i>Nucl. Data Tables A10</i>, 205 (1971); Log-<i>f</i> Tables for Beta Decay</p> | <p>71St47 R. M. Steffen—<i>Angular Distributions and Correlations of Radiation Emitted from Oriented Nuclei</i>; Report LA-4565-MS, Los Alamos Scientific Laboratory (1971)</p> <p>71St48 R. M. Steffen—<i>Proc. Int. Conf. Angular Correlations in Nuclear Disintegration</i>, Delft, Netherlands (1970), H. van Krugten, B. van Nooijen, Eds., Wolters-Noordhoff Publ., Groningen, p. 1 (1971); Angular Distributions and Correlations of Nuclear Radiations in Nuclear Spectroscopy</p> <p>71Ta32 W. H. Taylor, B. Singh, F. S. Prato, R. McPherson—<i>Nucl. Data Tables A9</i>, No. 1, 1 (1971); A Tabulation of Gamma-Gamma Directional-Correlation Coefficients</p> <p>72An20 I.V. Anicin, R.B. Vukanovic, A.H. Kukoc—<i>Nucl. Instrum. Methods</i> 103, 395 (1972); The New Instrum. of 1-3 Directional Correlations with Mixed Unobserved Transitions</p> <p>72Bb16 W. Bambynek, B. Crasemann, R. W. Fink, H.-U. Freund, H. Mark, C. D. Swift, R. E. Price, P. Venugopala Rao—<i>Rev. Mod. Phys.</i> 44, 716 (1972); X-Ray Fluorescence Yields, Auger, and Coster-Kroni Transition Probabilities</p> <p>72El21 Y. A. Ellis, M. R. Schmorak—<i>Nucl. Data Sheets B8</i>, 345 (1972); Survey of Nuclear Structure systematics for A ≥ 229</p> <p>72Tr09 V. F. Trusov—<i>Nucl. Data Tables</i> 10, 477 (1972); Energy Transitions</p> <p>76Ba63 I. M. Band, M. B. Trzhaskovskaya, M. A. Listengarten—<i>Nucl. Data Tables</i> 18, 433 (1976); Internal Conversion Coefficients for Atomic Numbers Z ≤ 30</p> <p>78Po08 F. T. Porter and M. S. Freedman—<i>J. Phys. Chem. Ref. Data</i> 7, 1267 (1978); Recommended Atomic Electron Binding Energies, 1s to 6p_{3/2} for Heavy Elements, Z=84 to 103</p> <p>79Ah01 I. Ahmad—<i>Z. Phys.</i> A290, 1 (1979); Precision measurement of K-Shell Fluorescence Yields in</p> |
|--|---|

ORGANIZATION OF MATERIAL

Within each A chain, information is collected by isotope and arranged in order of increasing Z. For each isotope, ^AZ, the arrangement of material and conventions for inclusion in tables are described below. Uncertainties are shown whenever available.

1. Adopted levels in ^AZ — All adopted level properties are shown for each level, together with explanatory comments.
2. Adopted γ radiations in ^AZ.
3. Radiations from ^AZ decay — All adopted radiation properties are shown for each mode of decay of the ^AZ ground state. Properties of α , β^- , AND $\epsilon + \beta^+$ radiations precede γ -radiation properties.
4. Radiations from ^AZ isomeric-state decay — Decay of each isomeric state is presented according to the same conventions as for the ground-state decay.

5. Levels and γ rays in ^AZ from decay[†] — Decays are ordered by A, Z of the parent.
 - a) Table of levels deduced from the decay.
 - b) Table of γ rays observed in the decay unless given under its respective parent according to 3 or 4 above.
6. Levels and γ rays in ^AZ from nuclear reactions[†] — Reactions are ordered by A, Z of the target, then by A, Z of the incident nucleus. A heading is given for each reaction.
 - a) Table of levels deduced from the reaction.
 - b) Table of γ rays observed in the reaction.

[†] The detailed results are tabulated only when the information is not adequately summarized on the drawings.

SUMMARY OF BASES FOR SPIN AND PARITY ASSIGNMENTS -continued

PROPOSITIONS ON WHICH STRONG ARGUMENTS ARE BASED

Ground States

1. The ground state of an even-even nucleus has $J^\pi=0^+$.
2. Spin determinations by such techniques as atomic-beam resonance, paramagnetic resonance, electron-spin resonance, and optical spectroscopy give correct values.

Gamma Transitions

3. The agreement of the measured value of a single conversion coefficient with the theoretical value for a multipolarity which is well separated from the value for any other multipolarity determines the transition multipolarity.
4. In all other cases if there is no other evidence for multipolarity, agreement of two or more measured conversion coefficients or ratios with theoretical values is necessary in order to establish the multipolarities of a transition and its mixing ratio.
5. Since E0 transitions can proceed only by conversion or pair production, pure E0 is ruled out if photons are observed.
6. Recommended upper limits for γ -ray strengths (Γ_γ/Γ_w , Γ_w -Weisskopf estimate) for various A-values are given below

$\Gamma_\gamma/\Gamma_w(\text{Upper Limit})$

Character*	A=6-44 ^{a§}	A=45-150 ^{a,b}	A>150 ^c
E1 (IV)	0.3 [#]	0.01	0.01
E2 (IS)	100	300	1000
E3	100	100	100
E4	100	100 [†]	
M1 (IV)	10	3	2
M2 (IV)	3	1	1
M3 (IV)	10	10	10
M4		30	10

* 'IV' and 'IS' stand for isovector and isoscalar.

† $\Gamma_\gamma\Gamma_w(\text{Upper Limit})=30$ for A=90-150

$\Gamma_\gamma\Gamma_w(\text{Upper Limit})=0.1$ for A=21-44

§ $\Gamma_\gamma\Gamma_w(\text{Upper Limit})=0.003$ for E1 (IS), 10 for E2 (IV), 0.03 for M1 (IS), 0.1 for M2 (IS)

^a From 79EnO4

^b From 81EnO6

^c Deduced from ENSDF by M. J. Martin

Beta Transitions§

- 7a. If $3.6 < \log ft < 5.9$, the transition is allowed: $\Delta J=0$ or 1, $\Delta\pi=+$ (no change in parity). For the mass region around Z=82, the upper limit should be lowered to 5.1.

See " *β -Decay Rate Probabilities*" on page App-F-iv. Note that $\log f^l u_t = \log ft + 1.079$.

* ($\log ft < 7.4$)

† ($\log ft \geq 7.4$)

- 7b. If $3.6 < \log ft < 6.4$, the transition is not $0^+ \rightarrow 0^+$. Superaligned ($\Delta T=0$) $0^+ \rightarrow 0^+$ transitions have $\log ft$ in the range 3.48 to 3.50. Isospin forbidden ($\Delta T=1$). $0^+ \rightarrow 0^+$ transitions have $\log ft > 6.4$.

8. If $\log f^l u_t < 6.5$, * $\Delta J=0,1$; $\Delta\pi=\pm$.

9. If $\log ft < 11.0$, $\Delta J=0,1$, $\Delta\pi=\pm$ or $\Delta J=2$, $\Delta\pi=-$

10. If $\log ft < 12.8$, $\Delta J=0,1,2$; $\Delta\pi=\pm$.

11. If $\log f^l u_t \geq 8.5^\dagger$ and if the Fermi plot has the curvature corresponding to a shape factor (p^2+q^2), then the transition is first-forbidden unique ($\Delta J=2$, $\Delta\pi=-$).

$\gamma\gamma$ Directional Correlation

$$W(\theta) = \sum_{k=\text{even}} A_k P_k(\cos \theta)$$

12. If a gamma-gamma directional-correlation experiment yields $A_2 \approx +0.36$ and $A_4 \approx +1.1$, then the spin sequence is $0 \rightarrow 2 \rightarrow 0$.

13. Results of $\gamma\gamma(\theta)$ are strong evidence for excluding spin sequences for which the theoretical A_2 or A_4 falls well outside the experimental range.

$\beta\gamma$ Directional Correlation

$$W(\theta) = \sum_{k=\text{even}} A_k(\beta) A_k(\gamma) P_k(\cos \theta).$$

14. If $|A_2(\beta)| \geq 0.1$ ($A_4=0$), the transition is not allowed. The converse is not true.

15. If $A_4(\beta) \neq 0$, the transition is neither allowed nor first-forbidden.

16. If $A_4(\beta)=0$, the transition is allowed or first-forbidden.

$\beta\gamma$ Polarization Correlation

$$P(\theta) = \frac{\sum_{k=\text{odd}} A_k(\beta) A_k(\gamma) P_k(\cos \theta)}{W(\theta)}$$

17. In allowed transitions,

$$\beta^- \quad A_1(\beta) < 0 \text{ if } J_i = J_f$$

$$\beta^+ \quad A_1(\beta) > 0 \text{ if } J_i = J_f$$

$$\beta^- \quad A_1(\beta) \leq 0 \text{ if } J_i = J_f + 1$$

$$A_1(\beta) < 0 \text{ if } J_i = J_f - 1$$

$$\beta^+ \quad A_1(\beta) \leq 0 \text{ if } J_i = J_f + 1$$

$$A_1(\beta) > 0 \text{ if } J_i = J_f - 1$$

18. If $A_3(\beta) \neq 0$, the β -transition is not allowed. The converse is not always true..

SUMMARY OF BASES FOR SPIN AND PARITY ASSIGNMENTS -continued

Reactions

19. Low-energy Coulomb excitation is predominantly E2 excitation.

20. Coulomb excitation determines J^π if the excitation probability agrees with the calculated values of Alder et al., Kgl. . Danske Videnskab. Selskab, Mat.-Fys. Medd. 32, No. 8 (1960).

21. The spin of the compound nuclear state resulting from thermal-neutron capture is equal to the spin of the target nucleus plus or minus 1/2.

22. Primary γ 's from neutron capture are E1, M1, E2, or M1+E2.

23. If the angular distribution in a single-nucleon transfer reaction can be fitted with a unique l -value, the spin of the final state J_f is related to the spin of the initial state J_i by

$$\vec{J}_f = \vec{J}_i + \vec{l} + \frac{1}{2}$$

with parity change if l is odd.

24. For $Z \lesssim 50$ and $Z \approx 82$, if the vector analyzing power for a single-nucleon transfer reaction shows a clear preference between $J=l+1/2$ and $J=l-1/2$ and if the l -value is known, then the J -value is determined.

The limitation in the regions of applicability results from a lack of measurements in other regions rather than an expected or observed violation.

25. If the angular distribution can be fitted with a unique L -value the J^π of the final state is related to the J^π of the initial state by

$$\vec{J}_f = \vec{J}_i + \vec{L}, \quad \pi_f \pi_i = (-1)^L,$$

for the following cases:

- A strong group observed in (p,t), (t,p), and (3He,n) reactions (strong groups are assumed to result from two identical nucleons transferred in a relative s-state)
- A strong group observed in the α -particle reaction (${}^6\text{Li,d}$).
- (e,e') and (α,α') inelastic scattering.

26. In reactions with $J^\pi=0^+$ target, projectile, and ejectile, if the yield of a group at 0° or 180° is

- non-zero, the parity of the final state is $(-1)^{J_f}$
- zero at several uncorrelated energies, the parity of the final state is $(-1)^{J_f+1}$

In reactions with a polarized $J^\pi=1$ projectile in the $m=0$ substate, with $J^\pi=0^+$ ejectile and target, if the yield of a group at 0° or 180° is

- non-zero, the parity of the final state is $(-1)^{J_f+1}$
- zero at several uncorrelated energies, the parity of the final state is $(-1)^{J_f}$.

§ See 73 Ra 10

Magnetic Moments

27. In nonrotational regions, if the known spin and magnetic moment of a level lead to an expected shell-model state by use of the Schmidt diagram, the level parity is determined.

28. In rotational regions, if the known spin and magnetic moment of a level agree with theoretical values for only one expected Nilsson level, the level parity is determined.

Deformed Region—Band Structure

Symbol	Evidence
D	Data from β - or γ -transitions suggest $J^\pi = 2^+$
x	The level is Coulomb excited
y	The level energy is given by the rotational level-energy formula
z,z'	The inertial parameter, decoupling parameters, fits the local trend

In terms of the above types of evidence, the following rules can be stated for nuclei in the rotational regions. $90 \leq N \leq 112$ and $Z > 87$.

29. For the first excited state in an even-even nucleus, x or Dz is sufficient to assign a spin and parity of 2^+ .

30. A level may be assigned to the ground-state rotational band on evidence of type xz or yz, provided the ground-state spin is well known.

31. A level may be assigned to a rotational band on evidence of type yz, provided at least one member of the band has well-known spin.

32. For $K=1/2$ bands, yz' is sufficient evidence to assign spins to the levels.

Alpha Decay

33. In odd-A nuclei, levels connected by α -transitions have the same spin and parity if the hindrance factor is less than 4.

34. For α -decay between two states, one of which has $J=0$, the parity change is given by $\Delta\pi = (-1)^{\Delta J}$.

SUMMARY OF BASES FOR SPIN AND PARITY ASSIGNMENTS -continued

PROPOSITIONS ON WHICH WEAK ARGUMENTS ARE BASED

1. In cases where gammas of one multipolarity "cluster" in one time region in the half-life vs. energy plot, as is true for M4's, other γ 's whose half-lives fall in this cluster may be assigned the corresponding multipolarity.

2. In cases where a cluster of two multipolarities occupy one time region, e.g. M1 and E2, a new gamma of which the half-life falls in this region may be assigned one of the two multipolarities or a mixture of the two.

3. Whenever $\Delta J \geq 2$, an appreciable part of the Gamma transition proceeds by the lowest possible multipole order. This statement is based on the scarcity or counter-examples and the observation that few E2 γ 's are as slow as M3's, few M2's as slow as B3's, etc.

4. Low-lying states of odd-A nuclei have shell-model spins and parities except in the regions where deformations appear. This argument is much stronger when supported by expected cross-section strengths (C^2S) in single-nucleon transfer reactions. It is recognized that some shell model predictions are stronger than others. For example, the shell model would mildly deny that the ground-state J^π of the 39th proton be $3/2^-$ but emphatically deny its being $3/2^+$. However, we have not included this distinction here and consider that all shell-model arguments are weak.

5. The spin and parity of a parent state may be inferred from the measured properties of its assumed isobaric analog resonance, and vice versa.

6. In regions of nuclear deformation, the Nilsson model can be used to limit the possible spins and parities.

7. Statements similar to 4 and 6 based on other models.

8. Statements based on interpolation or extrapolation of regional trends.

9. All statements connected with the nonobservation of expected transitions.

REFERENCES

- 73Ra10 S. Raman N. B. Gove, *Phys. Rev. C* 7, 1995 (1973); Rules for Spin and Parity Assignments Based on Log ft Values.
- 79 En04 P. M. Endt, *At.Data Nucl.Data Tables* 23, 547 (1979); Strengths of Gamma-Ray Transitions in A = 45-90 Nuclei.

81En06

P. M. Endt, *At.Data Nucl.Data Tables* 26, 47 (1981); Strengths of Gamma-Ray Transitions in A = 91-150 Nuclei.

CONVENTIONS USED IN NUCLEAR DATA SHEETS

Units

- Energies keV
- Cross Sections barns
- Magnetic dipole moments nuclear magnetons
- Electric quadrupole moments .. barns

Uncertainties ("Errors") The uncertainty in any number is given one space after the number itself:

- 4.623 3 means 4.623 ± 0.003
- 4.6 12 means 4.6 ± 1.2
- 5.4×10^3 2 means 5400 ± 200
- 4.2 +8-10 means $4.2^{+0.8}_{-1.0}$
- 4.2 +8-10 means $-(4.2 + 10-8) = -4.2^{+0.8}_{-1.0}$

? Question marks given after the energy value of a level or a radiation represent doubt as to the existence of that level or the radiation. A "?" given after the $T_{1/2}$ value indicates that the assignment of that half-life to the given level is not certain.

() Parentheses have the following interpretation for different quantities in the tabular data:

Quantity	Meaning of Parentheses
J^π	J^π based upon weak arguments
L, Mult.	Probable but not definite
Other	Value deduced (not directly measured) or taken from other experiment(s).

Examples:

- $J^\pi=(1/2,3/2)^-$
Weak arguments limit the spin to 1/2 or 3/2. Strong arguments indicate negative parity.
- $J^\pi=4 (+)$
Strong arguments show the spin is 4; weak arguments suggest positive parity.
- Mult.=M1(+E2)
Multipolarity is M1 with possible admixture of E2.



[] Brackets

- 7/2-[514] Nilsson asymptotic quantum numbers,
- $K^\pi[N n_Z \Lambda]$
- Assumed quantity, e.g., [M1+E2], [7/2+]

XREF (Cross Reference)

The cross reference symbols (characters 'A' to 'O') for the various experiments in which a level is seen are sometimes given in the adopted-levels table. When an adopted level may correspond to more than one level in a given experiment the flag for that experiment is given in lower case. In cases of ambiguity the energy for the corresponding level is given as a comment.

Level Scheme Symbols

- Level of which the existence is not well-established
- Adopted position of neutron- or proton-separation energy
- Well-established level
- Ground-state or long-lived isomeric level
- ↓ Radiation of which the position on the level scheme is not well-established
- ⋮ An expected strong transition that has not been observed
-  Beta, gamma, and alpha ray placement is consistent with coincidence measurements
-  4380 0.4 7.5
4380-keV α -transition with intensity 0.4% and hindrance factor (HF) = 7.5
-  Decay modes of a level
-  404 4 8.5
 β^- -transition with endpoint energy, 404 keV measured directly; intensity, 4% of decays, usually found indirectly (from γ -intensities, a more accurate method if there are several β -groups); $\log ft=8.5$
-  119 80 6.3
 ϵ_L -transition with energy determined directly from endpoint of γ -continuum (to which is added K-shell electron binding energy) or from L-, K-capture ratio; with intensity 80%, usually determined indirectly (from γ -intensities); $\log ft=6.3$
- ↓ 665 25
 γ -transition with energy 665 keV and intensity 25 in the units specified in the drawing
- % Number per 100 disintegration of a nuclear level

Nuclear Data Sheets Symbols and Abbreviations

A	mass number*, $A=Z+N$	IT	isomeric transition
A_2, A_4	coefficients of Legendre polynomials in angular-correlation or angular-distribution measurement	J	total angular momentum quantum number*
av	average	K	projection of nuclear angular momentum J on nuclear symmetry axis
B(EL), B(ML)	reduced EL, ML transition probability in $e^2 \cdot (\text{barn})^L, \mu_N^2 \cdot (\text{barn})^{L-1}$	K, L, M	K-, L- M-shell internal conversion
calc, CA	calculated, calculation	K/L	K-, L-conversion electron ratio
CCBA	coupled-channel Born approximation	L	(1)orbital angular momentum quantum number*, (2)multipolarity
c e	conversion electron	L(n), L(p)	L-transfer in neutron, proton transfer reaction
chem	chemical separation	min	minute
circ	circular	M+	$M+N+O+\dots$
c.m.	center of mass	M1, M2, ML	magnetic dipole, quadrupole, 2^L -pole
coef	coefficient	mag spect	magnetic spectrometer
coin	coincidence	max	maximum
Coul. ex.	Coulomb excitation	Moss	Mossbauer effect
CP	circular polarization	ms	(1)mass spectrometer, (2)millisecond
cryst	crystal diffraction spectrometer	mult	multipolarity/character
C^2S, C^2S'	one- nucleon spectroscopic strength for pickup, stripping reactions	N	neutron number*, $N=A-Z$
d	day	NMR, NQR	nuclear magnetic, quadrupole resonance
D	dipole	norm	normalization
DSA	Doppler shift attenuation	PAC	perturbed angular correlation
DWBA	distorted-wave Born approximation	pc	proportional counter
DWIA	distorted-wave impulse approximation	$p, \gamma(\theta)$	angular distribution of γ -rays with respect to a proton beam
E	energy	$p, \gamma(t)$	time distribution of photons with respect to a pulsed proton beam
$E(\epsilon)$	energy of electron-capture transition (endpoint of γ -continuum + K-electron separation energy of daughter)	pol	polarized, polarization
E1, E2, EL	electric dipole, quadrupole, 2^L -pole	priv comm	private communication
excit	excitation function	PWBA	plane-wave Born approximation
expt	experiment, experimental	Q	(1) reaction energy*, (2) disintegration energy*, (3) quadrupole moment*, in units of barns, (4) quadrupole
F	fission	$Q(\epsilon)$	total disintegration energy in ϵ decay
F-K	Fermi-Kurie (plot)	$Q(\beta^-)$	total disintegration energy in β^- decay
FWHM	energy resolution, full width at half maximum	$Q(\alpha)$	total disintegration energy in α decay $E(\alpha) + E(\text{recoil})$
g	gyromagnetic ratio*	R	$r_0 A^{1/3}$, nuclear radius*
GDR	giant dipole resonance	RDM	recoil distance measurement
GQR	giant quadrupole resonance	RUL	recommended upper limit for γ -ray strength
g.s.	ground state	rel	relative
h	hour	res	resonance
H	magnetic field	s	second
HF	hindrance factor	S	spectroscopic factor
hfs	hyperfine structure	S'	$[(2J_f + 1) / 2J_i + 1]S$
HI	heavy ion	S(n) or S_n'	energy necessary to separate a neutron, proton from nucleus
I	intensity	S(p) or S_p ,	
IAR	isobaric analog resonance	scatt	scattering
IAS	isobaric analog state	scin	scintillation counter
IBS	internal brehmsstrahlung spectrum	semi	semiconductor detector
IMPAC	ion implantation perturbed angular correlation technique	SF	spontaneous fission
inel	inelastic	spall	spallation
ion chem	chemical separation by ion exchange	sr	steradian

APPENDIX F-5

Nuclear Data Sheets Symbols and Abbreviations (cont'd)

syst, SY	systematics								
t	triton			δ	ratio of reduced matrix elements of (L+1)- to L-pole radiation with sign convention of Krane and Steffen, <i>Phys.Rev. C2, 724 (1970)</i>				
T	(1)isobaric spin, (2)temperature			ε	electron capture				
T _Z	Z-Component of isobaric spin, (N-Z)/2			$\varepsilon_K, \varepsilon_L, \varepsilon_M$	electron capture from K-, L-, M--shell				
T _{1/2}	half-life*			$\varepsilon(\gamma)B(E2),$	partial B(E2) for pphoton,				
th	thermal			$\varepsilon(ce)B(E2)$	conversion electron detection				
thresh	threshold			θ	indicates angular dependence				
tof	time-of-flight measurement			λ	(1)projection of particle angular momentum on nuclear symmetry axis, (2)radiation type, e.g., M1, M2...				
vib	vibrational			μ	magnetic moment of particle*, given in nuclear magnetons (μ_n)				
W.u.	Weisskopf single-particle transition speed			ν	neutron shell-model configuration				
y	year			π	parity, proton shell-model configuration				
Z	atomic number*, Z=A-N			σ	cross section*				
α	total γ -ray internal conversion coefficient $N(ce)/N(\gamma)^*$			$\Sigma(\gamma\gamma)$	coincidence summing of γ -rays				
$\alpha(K), \alpha(L)$	γ -ray internal conversion coefficient for electrons ejected from the K-, L--shell			$\omega(K), \omega(L)$	K, average-L fluorescence yield				
$\alpha\gamma, \beta\gamma, \gamma\gamma$	coincidences of α 's and γ 's, β 's and γ 's, and γ 's and γ 's			% α	percent α branching from level				
$\alpha\gamma(\theta, H, t),$ $\beta\gamma(\theta, H, t),$ $\gamma\gamma(\theta, H, t)$	$\alpha\gamma$ -, $\beta\gamma$ -, $\gamma\gamma$ - coincidences as functions of angle, magnetic field, time			% β -	percent β - branching from level				
$\beta_2, \beta_3, \beta_L$	quadrupole, octupole, 2 ^L -pole nuclear deformation parameter			% β +	percent β + branching from level				
$\beta\gamma(pol),$ $\gamma\gamma(pol)$	polarization correlation of γ 's in coincidence with β 's, γ 's			% ε	percent ε branching from level				
$\Gamma, \Gamma(\gamma), \Gamma(n)$	level width*, partial width for γ -, n-emission			%IT	percent ($\gamma+ce$) branching from level				
$\gamma(\theta, H, T)$	γ -intensity as function of angle,			%SF	percent spontaneous fission from level				
γ^\pm	annihilation radiation			$\langle r^2 \rangle$	root-mean-square of nuclear radius				
Prefixes*				Symbols for Particles and Quanta *					
T	tera	(=10 ¹²)	m	milli	(=10 ⁻³)	n	neutron	π	pion
G	giga	(=10 ⁹)	μ	micro	(=10 ⁻⁶)	p	proton	μ	muon
M	mega	(=10 ⁶)	n	nano	(=10 ⁻⁹)	d	deuteron	e	electron
k	kilo	(=10 ³)	p	pico	(=10 ⁻¹²)	t	triton	ν	neutrino
c	centi	(=10 ⁻²)	f	femto	(=10 ⁻¹⁵)	α	α -particle	γ	photon
			a	atto	(=10 ⁻¹⁸)				

*Recommended by Commission on Symbols, Units, and Nomenclature of International Union of Pure and Applied Physics